

I. FINETTI **Universität Hamburg**
Institut für Geophysik
- Bibliothek -
 Bundesstrasse 55
 D-2000 HAMBURG 13

27. Juni 1983

STRUCTURE, STRATIGRAPHY AND EVOLUTION OF CENTRAL MEDITERRANEAN

Summary. The purpose of this paper is to outline the regional structure, stratigraphy and evolution of the Pelagian and Ionian Seas. Information, descriptions, remarks, ideas and conclusions here reported are based on a great amount of new geological, geophysical and drilling exploration data, and on the numerous most informative published scientific papers.

The studied area has been schematically subdivided into geological provinces according to their relevant regional characters. Every province is described in a synthetic manner regarding stratigraphy, structural setting, main geodynamical phases, which have occurred and associated regional volcanic activities.

Four principal extensional phases are recognized from Permo-Triassic to Quaternary. The first one, active during Middle-Upper Triassic, produced a continental rifting with prominent effects in the Gabes-Tripoli-Misurata basin, Steppenosa Trough (Sicily-Malta platform area), Sicily-Malta-Medina Mounts and Medina Bank areas, and in the Ionian Sea. The second extensional phase occurred in Middle-Jurassic and opened the Ionian Sea. The Sicily-Malta escarpment is affected by an extensional mega-fault system that connects the thick continental crust of the Pelagian Sea to the paleo-oceanic crust of the Ionian abyssal basin.

The Pelagian Sea and Sirte rise (excluding the Upper Sirte Slope, where sedimentation commenced in the Upper Cretaceous), are constituted by a more or less thick sedimentary sequence that includes depositions from Triassic to Quaternary. The Sirte Rise, as well as other geological provinces of the studied area, was greatly stretched during the third extensional phase of the Middle Upper Cretaceous.

The Ionian abyssal basin is interpreted as a paleo-oceanic crust which has been continuously covered by deep water sediments from the Middle Jurassic to Quaternary. In the central part of the Ionian, where there exists the maximum Bouguer anomaly of the Mediterranean, the sedimentary sequence as well as the lower crust, are considerably thinner.

The last main extensional phase occurred from the Middle-Upper Miocene to Quaternary and affected the greater part of the Pelagian and Ionian Seas, but in particular the areas of Malta-Pantelleria-Linosa, Medina Grabens and the Ionian Sea. Malta Islands are structurally an horst which emerged in the Pelagian Sea during late Miocene-Lower Pliocene. The rifting process of the Pelagian Sea involves the entire median area known as the Sicily Channel with numerous extensional faults, block collapsing, tilting and widespread volcanic activity.

Received August 1, 1982

1. Introduction

After more than two decades of intense geological and geophysical exploration activities, remarkable progress has been made in the knowledge of structure and stratigraphy of the Mediterranean. This does not mean that all aspects of the basin are now fully understood. But it is today possible to propose schemes of geodynamical evolution supported by more data. In addition to the scientific investigation, a consistent contribution has been made also by the oil exploration activity on the

La responsabilità scientifica degli scritti rimane agli autori

Fotocomposizione: Studio Gamma - Trieste
 Stampa: Ricci - Trieste
 Autorizzazione del Tribunale di Trieste, n. 242 del 17 settembre 1960

Istituto di Geodesia e Geosifica, Università di Trieste

continental shelf areas. Drilling results of the JOIDES project in deep waters are also very important, especially for the study of the Messinian and Plio-Quaternary series.

In the present paper, the Author outlines the regional stratigraphic and structural conditions of the Central Mediterranean (Pelagian and Ionian Seas) mainly resulting from interpretation of the seismic exploration conducted by OGS, or published, or available. Moreover some major extensional phases, associated volcanic activities, and subsidences are recognized and described.

Careful analyses of the great amount of collected data allow the existence of an old oceanic crust in the abyssal plain of Ionian Sea to be deduced. The geological interval of time of crustal opening and evolution of such a basin are also proposed.

The investigation is extended also to the southern Apulian platform area which is considered as a plate separated from the African megaplate during the crustal opening of the Ionian Sea.

An important contribution to the understanding of the geodynamical movements of the studied area is given by the reconstruction of the volcanic activities which have occurred during various phases from the Triassic to the present time.

Significant contributions to the enhancement of knowledge were furnished by the drilling exploration activities of Glomar Challenger (D.S.D.P.; Legs 13 and 42 a); Ryan et al., (1973); Hsü et al., (1978).

It has been clear for several years that the Ionian Sea is a key area for the observation of important geological events, determinant for the understanding of the evolution of the Central-Eastern Mediterranean. For this reason many Authors were attracted to the exploration of this basin.

After the discoveries of the relevant data in support of the plate tectonics theory in the Atlantic ocean and its margins, the first general attempt to introduce a plate tectonic model in the whole Mediterranean, to explain the evolution of this basin, was carried out by Dewey et al., (1973). The Authors of this paper were of the opinion not only that the history of the Mediterranean and Alpine system is basically associated with the opening of the Atlantic and relative plate movements between the Americas and Europe-Africa, and between Africa and Europe, but also that the fundamental plate tectonic models, developed at that time, were valid for the Mediterranean area. It is to be said that at the time of publication of this paper the exploration and knowledge of the Mediterranean was much more scarce than some years later. For this reason the Authors had not the opportunity to take into consideration some important data and aspects observed by successive explorations, but they had the merit to believe in the plate tectonic models for the Mediterranean, even if at that time the Authors favourable to such an interpretative approach were not numerous.

Boccaletti et al., (1972; 1974; 1976) applied progressively the interpretation model of Marginal and Back-arc basin to the Ligurian Sea and Apennines and, at the end to the entire Mediterranean area.

Biju-Duval et al., (1976; 1977) proposed a plate tectonic model based on the kinematic evolution of the African and European plates given by Pitman and Talwani (1972), and using the rotation suggested by Francheteau (1973), for the relationships between Africa and North America. This interesting model takes into account a very consistent number of observations and considers a more limited number of plates than Dewey et al., (1973).

Laubscher (1971; 1973; 1974) and Laubscher and Bernoulli (1977), examining the Central-Western Alps, the Dinarides and Hellenides, propose plate tectonic models having some original modifications with respect to the classic schemes. These Authors emphasize the palinspastic reconstructions and consider gravity as the main driving force in the subduction process. The model which postulates a southern origin of




Fig. 1 — Studied Area of Central Mediterranean and Scheme of Geological Subdivision.




Fig. 2 — Bouguer Gravity Map of Central Mediterranean.




Fig. 3 — Magnetic Total Intensity Map of Central Mediterranean.

Antalya-Cyprus Hatay ophiolites and the opening of the Eastern Mediterranean in Mesozoic (Laubscher et al., 1977) is similar in some aspects to that of Biju-Duval et al., (1976).

Based on a deep knowledge of stratigraphic sequences of the Southern Apennines, and assuming several concepts already presented in Laubscher et al., Scandone et al., (1977) propose an interesting geodynamic model of the Central Mediterranean area.

Carey (1976) presents a global geodynamic evolution based on his expanding theory published in 1953. The Author interprets the evolution of Tethys as a consequence of torsional megadeformations, with rotation of blocks.

An original approach to the evolution of the earth's crust and in particular of the Alpine System in the Mediterranean is proposed by Tapponier P. (1977). Following this theory and interpretation, the African promontories of the Italian and Arabian peninsulas played an important role in determining the tectonic conditions in the collision area with stable Europe.

Channel, D'Argenio and Horvath (1979) reconstruct the paleotectonic evolution of the "Adria" plate, considered as a promontory of the African one. The periadriatic orogenic systems are examined and the ideas expressed are supported by several consistent arguments, and new data or data available from careful consultation of the whole literature. Also Semenza (1977), studying in detail the Southern Alps, arrives at the conclusion that the Adriatic (Italo-Austro-Dinaric plate) is substantially a promontory of Africa.

Bosellini (1973) discussed a geodynamic model of the Southern Alps in Jurassic and Cretaceous. Castellarin and Vai (1981) remark on the importance of Hercynian tectonics in the Alpine system. Castellarin, Colacicchi and Praturlon (1978) reconstruct the extensional tectonics along with the important geological area of the Ancona-Anzio line.

Many other important papers have contributed to the progress in the knowledge of the Mediterranean, and only the limit of space does not allow the enumeration of other relevant research works to be continued.

2. Regional structural and stratigraphic conditions of Pelagian and Ionian Seas

All reconstructions of the regional structural and stratigraphic conditions are mainly based on seismic explorations and on remarkable amounts of available data of oil exploration along the continental margins. The most complete regional seismic exploration which covers the whole Mediterranean is that carried out by the OGS of Trieste, and partially published by Finetti and Morelli (1972; 1973) and Finetti (1976, 1981).

Important seismic information on deep water areas of the Mediterranean are given also by the exploration activity of IFP, CNEXO and others: Mauffret et al., (1973); Biju-Duval et al., (1974); Mulder (1973).

These data, together with the drilling results of the JOIDES project (Legs 13 and 42 a) constitute the most useful information for the outline of the existing geological conditions of deep water areas of the Mediterranean.

Particular attention has been dedicated to the Mesozoic extensional geodynamics because of its importance for the reconstruction of the evolution of the area. While of the Pelagian and Western Ionian Sea the tectonics associated with the Mesozoic movements is still evident, in the Eastern Ionian Sea, on the contrary, the successive compressional movements have largely modified the previously existing tectonic setting. Only in a small area between the Calabrian and Hellenic Arcs it is still possible to observe the extended margin of Apulia (Figs 4 & 5).

From the stratigraphic and tectonic viewpoint it is possible and convenient to divide the studied areas into seven geological provinces (Fig. 1):

- Pelagian Sea
- Sirte rise
- Ionian abyssal basin
- Apulian platform
- Southern Adriatic basin
- Calabrian arc
- Hellenic arc

These provinces have individual characters which are remarkably different from each other. But in every province its own particular characteristics can be substantially observed over the whole area.

The Pelagian Sea is the most explored province of those studied in this paper, and for its varying characteristics it has been subdivided into seven different geological units (Fig. 1):

2.1 — Pelagian Sea

With this name the area between Sicily, Tunisia (Cape Bon), Tripolitania coast, the upper limit of the Sicily-Malta escarpment and the Medina bank to the east is intended. In literature this part of the Mediterranean is also called the "Strait or Channel of Sicily". We will limit the term "Channel of Sicily" only to the central area of the Pelagian Sea where the prominent rifting process exists and the larger grabens like those of Malta, Pantelleria, Linosa and Medina occur.

To illustrate and describe the varying geological and geophysical aspects of this Sea it is convenient, as above mentioned, to subdivide the area into the following main geological units (Fig. 1):

- Gages-Tripoli-Misurata basin
- Lampedusa-Medina plateaus
- Sicily Channel rifting area
- Caltanissetta basin
- Ragusa-Malta plateau
- Adventure plateau
- Sicily-Malta-Medina Mounts Escarpment

The last unit (Sicily-Malta-Medina Mounts escarpment) does not belong strictly to the Pelagian Sea. But to follow the relationships between this area and the Ionian abyssal basin, and to examine the remarkable difference between the steep escarpment of Sicily-Malta and that much more gentle of Sirte rise, it is better to give its description immediately after the units previously commented on.

2.1.1 — Gages-Tripoli-Misurata Basin

This geological unit extends from the Sfax-Gages area to offshore Misurata. It is limited to the south by the regional extensional fault system of Gafsa-Jeffara, and to the north by the Lampedusa-Medina Plateaus area.

From the tectonic viewpoint this basin is to be considered as the result of old extensional geodynamical movements which occurred in the Triassic and before in the Paleozoic. The two flanks, as well as the trough of the basin are affected by several extensional faults, some of which with remarkable displacements.

Extensional phases, successive to the first ones above mentioned, activated the previously existing faults, and created new ones. After the Paleozoic-Triassic stretching movements other important extensional geodynamic phases occurred in the Middle Jurassic and especially in the Middle-Upper Cretaceous. With this last one are associated remarkable igneous effusions of a basaltic type. Basaltic layers in the Middle-Upper Cretaceous have been found in several points by drilling exploration. These movements were accompanied by a consistent subsidence of the basin. Extensional and subsidence activities occurred also successively. In the Middle-Upper Miocene to Quaternary, one of the main extensional phases which affected not only and not particularly this area took place.

On seismic lines it is possible to identify many more faults than it is possible to indicate on a map on a regional scale like that of *Plates III* and *IV*.

Combining seismic stratigraphy and available borehole data it is possible to reconstruct the following average sequence (thicknesses of Lower Mesozoic in metres by conversion of traveltimes following the obtained seismic velocities):

- a) Plio-Quaternary (100-350)
- b) Messinian interval (0-150)
- c) Middle Miocene to Oligocene (1400-2200)
- d) Eocene-Paleocene (700-1000)
- e) Cretaceous (1200-2000)
- f) Jurassic (1500-2600)
- g) Permo-Triassic (Probably more than 3000)

Triassic salt walls are known to exist from boreholes made. Seismic evidence of Triassic salt walls, especially on the deeper part of the Gabes-Tripoli basin has been found. From seismic velocity functions it seems clear that Miocene-Oligocene is generally associated with relatively low values (2000 to 3000 m/sec) attributable to predominant shaly-marly sequences. Eocene and Mesozoic are frequently characterized by higher velocities probably corresponding to carbonate intervals, homogeneous and/or interbedded with shaly-marly layers.

2.1.2 — Lampedusa-Medina Plateaus

Along with the median area of the Pelagian Sea there exist two regional structural highs: that of Lampedusa and that of Melita-Medina banks. Even if these structures show several local variations with very gently undulations and many young normal faults with horst and grabens, schematically we can consider them, in a regional manner, as two large plateaus. Such features are limited to the south by the Gabes-Tripoli-Misurata basin, and to the north by the troughs of Linosa and Medina Grabens *Plate III*. While the Lampedusa Plateau is regionally an unique high, the Melita-Medina plateau is much more fragmented and constituted by different highs and lows affected by numerous faults.

The Lampedusa plateau is higher than the Melita-Medina one. To compare the height of the two structures the Lampedusa area has been assumed as that inside the isochrone of 1.0 sec, while the Medina Bank area has been assumed as that inside the isochrone of 1.6 sec (top Mesozoic, *Plate III*). So Lampedusa plateau on average results about 0.6 sec, or about 0.9 km higher.

Examining the geological evolution of these parts of the Pelagian Sea, it seems possible to say that the Lampedusa-Medina plateau is a regional high area that has existed for a long time: at least since the Cretaceous. The more recent movements of the Miocene-Quaternary extensional phase produced fracturation with creation of local uplifted blocks and grabens within the regional swell.

Both the Lampedusa plateau and the Melita-Medina plateau areas are affected by more or less consistent volcanic activities. The Medina Bank especially shows several prominent magnetic anomalies. Probably the Cretaceous phase is the most consistent and widely diffused one. But also the last extensional phase (Miocene to Quaternary) seems to have locally affected the area along with the major faults.

By interpolation and extrapolation on seismic lines of the conditions known in the Gabes-Tripoli-Misurata basin and on the Sicily-Malta plateau it is possible to predict the following average stratigraphic condition for the Lampedusa and Medina Bank areas (thickness in metres):

- 1) *Lampedusa plateau area*
 - a) Plio-Quaternary (0-350)
 - b) Messinian (0-50)
 - c) Middle Miocene to Oligocene (1200-1500)
 - d) Eocene-Paleocene (500-900)
 - e) Cretaceous (2200-3000)
 - f) Jurassic (1200-2200)
 - g) Permo-Triassic (More than 2000)
- 2) *Medina bank area*
 - a) Plio-Quaternary (0-250)
 - b) Messinian (0-50)
 - c) Middle Miocene to Oligocene (200-900)
 - d) Eocene-Paleocene (200-900)
 - e) Cretaceous (3000-5000)
 - f) Jurassic (2400-3000)
 - g) Permo-Triassic (More than 2000)

At the Lampedusa Island a Cainozoic-Mesozoic limestone sequence is outcropping and the structural uplift is limited to the north by a remarkable fault. The predominant trend is that parallel to the axis of the Pelagian Sea: NW-SE. At the Medina bank area the Tertiary sequence is generally thicker than on the Lampedusa Plateau. Mesozoic, following our seismic interpretation, seems substantially comparable in terms of thickness in the two considered plateaus.

2.1.3 — Sicily Channel Rifting Area

The morphological setting of the Pelagian Sea, for the most part, is relatively young and associated with geotectonic conditions.

The regional structural conditions can be assumed as substantially tabular on the fractured parts of the basin and everywhere before the "NQ" rifting phase. During the last main extensional phase of the Miocene-Quaternary a remarkable stretching and faulting activity affected the whole Pelagian Sea and in particular the area of the Sicily Channel in correspondence with the major grabens of Pantelleria, Linosa, Malta and Medina (*Plate III*). Before this geodynamical phase, a long period of relatively not important movements or quiet conditions occurred in the Pelagian Sea. During this interval of time the whole Pelagian Sea area was a basin where sedimentation took place with continuity on a sea bottom having moderate depth variations from place to place, with few limited exceptions, like for example the areas of Lampedusa and Lampione Islands. Malta Island, from the whole Middle Miocene back in time to at least the Cretaceous, did not exist.

The Middle-Miocene-Quaternary stretching and rifting activity produced here the most widely distributed deformation process with numerous normal faults, some having remarkable displacements (Figs 11, 12, 13). During this geodynamic activity the Pelagian sea assumed the present sea-bottom setting with the generation of major grabens like those of Malta, Pantelleria, Linosa, and Medina (Fig. 9). In the Upper Miocene-Early Pliocene, or about 10 m.y. ago, the block of the Maltese Islands emerged. Examining in detail the structural setting and evolution of the Malta graben and Maltese Islands it is possible to make some important remarks:

- a) Maltese Islands block is structurally a Horst (Fig. 11).
- b) The extensional and rifting movements of major grabens as well as that of Malta, are generally accompanied by tilting processes of the two uplifted blocks on both sides of the considered graben. Maltese Islands are generated by the tilting of the NE block of the Malta Graben.

Rifting movements of this "NQ" phase are also associated with a prominent volcanic activity which occurred in several parts of the Sicily Channel. The most important outcropping volcanic shows are those of the Pantelleria and Linosa Islands. But exploration activity and magnetic anomalies indicate that several other areas were affected by volcanism from the Miocene to the Quaternary. See for example Medina Graben (Fig. 9).

The stratigraphy of this area, following seismics, shows appreciable variations in terms of thickness and probably also of lithology. In fact it is possible to see horizontal seismic velocity changes along with equivalent time-stratigraphic intervals. In the following list the time stratigraphic intervals of the Malta area resulting from seismic exploration are given:

- a) Marine Plio-Quaternary and Messinian (Missing)
- b) Miocene to Oligocene (500-600)
- c) Eocene-Paleocene (700-1000)
- d) Cretaceous (1200-1500)
- e) Jurassic (1800-2200)
- f) Permo-Triassic (More than 3000)

Here, as well as on the whole area from Ragusa to Medina Mounts, the Middle Jurassic is affected by the most impressive volcanic activity (Phase "J"). More modest volcanic shows occur also in the Cretaceous extensional phase ("K"). The entire Mesozoic sequence of the Malta area furnishes high seismic velocities, in accordance with the existence of a carbonate sequence which is here much more continuous than in the Ragusa and the Medina bank areas.

2.1.4 — Caltanissetta Basin

Central Sicily is geologically characterized by a very thick Neogene basin known as "Caltanissetta Basin". The thickest stratigraphic sequence is that of Miocene (2000 to 4000 m) which terminates with the late Miocene evaporitic interval. This evaporitic succession in the Caltanissetta basin is constituted by evaporitic and terrigenous sediments cyclically alternated. Evaporitic deposits here are so typical that from the town of Messina, this chronological interval (Messinian) takes its name.

Since the Early Neogene the basin has been affected by a marked subsidence accompanied by high rate deposition. From the Middle Miocene to the Quaternary, both the gravitative sliding and the compressional geodynamics determined an exasperate tectonics with shearing planes, overthrusts and chaotic allochthonous conditions.

Movements continued also during the Quaternary as the existence of the Quater-

nary olistostrome in South Sicily and the Gulf of Gela (Plate III), indicates.

This basin extends in the Pelagian Sea towards SE in the area north of Malta. Stratigraphically, the area between Malta and Ragusa, from the Eocene to the Mesozoic is similar to that of Malta. But here also the Oligocene and a thicker Miocene, and thin Plio-Quaternary exist. More detailed stratigraphical and structural conditions are reported in Patacca et al., (1981).

2.1.5 — Ragusa-Malta Plateau

The area between Ragusa and Malta, limited to the East by the Sicily-Malta escarpment, is constituted by a very thick Mesozoic sequence covered by a Cainozoic progressively thinning from west to east and from south to north. In south-east Sicily, and near the limit of the escarpment, the Mesozoic outcrops, or is very shallow. The whole Ragusa-Malta area shows gentle undulation mostly trending NE-SW in the northern part. But also NW-SE less pronounced trends can be identified. Faults are not so numerous as in the area of the major grabens (Malta, Pantelleria, Linosa ad Medina grabens) and are in general older because they are frequently associated with the Mesozoic extensional phases ("J", "K"). In some cases the paleo-fault systems stopped their activity during the first generating extensional phase; in some others they were renewed by successive phases. Regionally, the Ragusa-Malta platform area can be considered as an asymmetric ridge trending N-S and affected by clear and remarkable tilting movements near the Sicily-Malta escarpment. Such tilting is associated with the main extensional phase which is here that of the Middle Jurassic ("J"). But also the Cretaceous ("K") and Neogene-Quaternary phase ("NQ") continued the uplifting and tilting movements of the block.

The Maltese Islands are constituted by small emerging areas of the uplifted block which limits to the North the Malta graben. Uplifting of this block and the emersion of the Maltese Islands are due to the tilting movements which is associated with the "NQ" rifting of the Sicily Channel area. The Maltese Islands block structurally is a horst (Fig. 11) from which the Islands emerged about 10 m.y. ago, at the end of Miocene.

The whole Ragusa-Malta platform area is affected by prominent volcanic activities which occurred at different regional stretching phases. Four main phases (Fig. 32) can be identified: Middle-Upper Triassic ("T"), Middle Jurassic ("J"), Upper Cretaceous ("K") and Neogene-Quaternary ("NQ").

In this area, all the phases produced intense activities with thick to very thick volcanic intervals. Particularly impressive is the Middle Jurassic phase. In fact, the relative interval is almost completely constituted by igneous rocks. But also the Middle-Upper Triassic (Patacca et al., 1981) shows a remarkable volcanic activity with many basaltic intervals in "Streppenosa" formation (Black Shale).

2.1.6 — Adventure Plateau

The area offshore of SW Sicily, known as "Adventure Plateau" is substantially flat at the sea bottom surface, but shows remarkable geodynamic movements and tectonization in subsurface. Both extensional and compressional movements, with predominance of the first one, affected the area. Extensional deformation and associated normal faulting is predominantly trending parallelly to the Sicily Channel (WNW-ESE). Compressional effects and reverse faulting, generally parallel to the Maghrebian trend (NE-SE), are much more modest than the extensional ones.

From SW Sicily toward Pantelleria Island the reflecting sequence shows a general progressive thickening of Mesozoic and Lower Tertiary. The Middle-Upper Miocene is characterized by a prograding sequence overlying a relatively thin Lower Miocene. Generally Jurassic is thin to very thin or missing.

Average time-stratigraphic sequence can be summarized as follows (thickness in metres):

- a) Plio-Quaternary (0-1000)
- b) Messinian (0-200)
- c) Middle-Upper Pre-Messinian Miocene (200-2000)
- d) Lower Miocene (0-800)
- e) Eocene-Oligocene-Paleocene (200-800)
- f) Cretaceous (600-1400)
- g) Jurassic (0-500)
- h) Triassic (> 3500)

The "NQ" extensional phase produced the most remarkable deformations with several volcanic shows, especially in the Southern part of the plateau.

2.1.7 — Sicily-Malta-Medina Mounts Escarpment

A very important geological meaning is assumed by the Sicily-Malta escarpment. Tectonically, it is affected by a normal regional mega-fault system that diminishes its total displacement from Sicily (Fig. 6) to the Medina Mounts (Fig. 7) about 1.5 second, two-way time, at Middle Jurassic layers.

South of the Ragusa Plateau (Fig. 6) there exists a complete thick Mesozoic, covered by a thin Eocene (about 200 m). Then a very big extensional fault system brings to the down block of the Ionian abyssal area with a slope directly constituted by the fault plane. It is evident, even from seismic characters, that up-block and down-block have very different Tertiary sequences: very thin and only of Eocene age the Upper block, and very thick and with a much more complete sequence the down one. Stratigraphy and tectonics of the down block are very indicative of the involved geodynamics. In fact it is possible to remark that the fault system commenced its activity in the Mesozoic and continued in the Tertiary. In particular during the Miocene a prominent foundering of the Ionian Sea as a consequence of the "NQ" extensional phase occurred.

The most important and significant difference between the up and down block of this area is not in the Tertiary, but very probably, in the Mesozoic and in the Crustal characters. After several years of geophysical, geological, and drilling exploration activities, studying all obtained data, the Author has arrived at the conclusion that the Ionian abyssal basin is constituted by a paleo-oceanic crust open during the Mesozoic. Arguments in support of such an interpretation are discussed later on in paragraph 2.3.

Structurally, the Medina Mounts are constituted by a complex fractured horsts system trending W-E in the western part (Fig. 37). Associated prominent magnetic anomalies (Fig. 3) in addition to seismic information (Fig. 7) indicate that this feature is affected by volcanism.

Examining more carefully the seismic exploration results it is possible to remark and/or deduce the following:

- a) The northern flank correlates very well, in terms of existing extensional faults and age of generating geodynamics with the Sicily-Malta escarpment. All geophysical data indicate that the northern escarpment of the Medina Mounts is the continuation of the Sicily Malta escarpment. In the Medina Mounts the total displacement of the involved extensional fault system is smaller than in the Sicily-Malta escarpment.
- b) The blocks forming the feature are differently foundered with respect to the Malta-Medina shallow platform area up to a depth of more than 2000 metres (Fig. 7). The foundering process of these blocks, very probably, commenced during the Cretaceous extensional phase ("K") where the whole Sirte Rise was greatly stretched. But the geodynamic phase that contributed the most is evidently that of




Fig. 1 — Example of Interpreted Seismic Reflection Line on Ionian Margin of Apulian Plate (Line MS-68; S.P. 1190-920). Evidence of extensional geodynamics with fragmentation and tilting.

APULIAN PLATFORM



LINE MS 69




Fig. 5 — Example of Interpreted Seismic Reflection Line on Ionian Margin of Apulian Plate (Line MS-69; S.P. 1730-1925). It shows the same geodynamical process of Figure 4.




Fig. 6 — Example of Interpreted Seismic Reflection Line on Sicily-Malta Escarpment (Line MS-26; S.P. 2147-1840).

It shows the passage from Ragusa-Malta Plateau area to the Ionian conditions. Evidence of old extensional geodynamics, with mega-fault system and crustal opening.

MEDINA MOUNTS




Fig. 7 — Example of Interpreted Seismic Reflection Line Across Medina Mounts Area (Line MS-21; S.P. 1940-1410). The Mega-Fault System between Medina Mounts and the Ionian Basin is the continuation of the Sicily-Malta Escarpment seen in Figure 6. Evidence of young volcanic activity associated with the "NQ" extensional phase.

LINE MS 14

MEDINA BANK




Fig. 8 — Example of Interpreted Seismic Reflection Line On Medina Bank and Medina Escarpment Area (Line MS-14; S.P. 4510-4110). It shows the continuity of the Mesozoic-Tertiary African Plate Sequence, not affected by faults in Medina Bank area. The extensional faults of Medina Escarpment were produced during the "K" and "NQ" stretching phase.

MEDINA GRABEN MIOCENE TO QUATERNARY RIFTING

LINE MS 14

N W

5625

5505

5405

MS-20



5195 SE




Fig. 9 — Example of Interpreted Seismic Reflection Line Across Medina Graben Area (Line MS-14; S.P. 5625-5195). Medina Graben is created during the "NQ" extensional geodynamics, as well as all major grabens of Central Pelagian Sea. Evidence of young volcanic activity (Plio-Quaternary) associated with prominent magnetic anomaly. Remarkable tilting process involving also deep layers and very probably lower crust.

S E EXTREMITY OF CALTANISSETTA BASIN

LINE MS 19

SW

480

RAGUSA
PLATEAU

NE




Fig. 10 — Example of Interpreted Seismic Reflection Line Across the South-East Extremity of Caltanissetta Basin (Line MS-19; S.P. 480-1).

LINE MS 19
SW 980

MALTA GRABEN

MALTA HORST



NE




Fig. 11 — Example of Interpreted Seismic Reflection Line Across Malta Horst and Malta Graben Area (Line MS-19; S.P. 980-480). It is the continuation of previous Figure, and the four Figures 10, 11, 12 and 13 show the entire line MS-19. In this Figure 11 it is evident that the structure of Malta corresponds to a horst. Malta Graben is affected by a prominent rifting process occurred in "NQ" extensional phase.

LINE MS 19
SW 1570

SICILY CHANNEL RIFTING AREA



NE




Fig. 12 — Example of Interpreted Seismic Reflection Line Across the Rifting Area of Central Pelagian Sea (Line MS-19; S.P. 1570-980). Intense faulting process with back-tilted blocks and some magmatic extrusions all associated with "NQ" stretching phase.




Fig. 13 — Example of Interpreted Seismic Reflection Line Across the Rifling Area of Central Pelagian Sea and the Tabular Area of Lampedusa Plateau (Line MS-19; S.P. 2060-1570). It shows the end of the rifted area and the much more quiet conditions of the Lampedusa Plateau.

- the Miocene-Quaternary ("NQ"). On the contrary during the crustal opening phase "J", this area was very probably uplifted by tilting movements.
- The Mesozoic blocks of the Medina Mounts are covered by different thicknesses of Tertiary from relatively thin to very thin. In some cases the Mesozoic outcrops.
 - The southern flank of the Medina Mounts is determined by extensional faults having in general a much more modest displacement (Fig. 7). The faults very probably commenced in the Middle-Upper Cretaceous (phase "K"), during the generalized stretching activity which affected the Sirte Rise. During the Miocene-Quaternary extensional phase, all the major existing fractures were newly activated and the displacements of the faults of the southern flank of the Medina Mounts are mostly due to these last geodynamical movements.

2.2 — Sirte Rise

With this term the whole Sirte margin area between the Sirte coast, the east limit of the Medina and Melita plateau and the Ionian abyssal basin is identified (Fig. 1 & Plate III).

The regional structure of this geological province can be schematically considered as a monocline affected by numerous extensional faults accompanied by tilted blocks. Assembling numerous exploration data, and extending the calibrated or reconstructed stratigraphic and structural information to the deep water of the Sirte Rise, it is possible to summarize the following remarks:

- In the area of the Gulf of Sirte, as is well known, on a Cambro-Ordovician or Hercynian basement, marine sedimentation commences in the Upper Cretaceous and continues until the Quaternary.
- Off the upper slope (Plate III) from about the isochrone 3.0 to 3.6 sec area, below the calibrated Upper Cretaceous sediments, a reflecting sequence, that can be correlated with the Lower Cretaceous to Triassic sequence known in the surrounding Pelagian Sea and Cyrenaica areas onlaps and progressively thickens seaward. Anyhow, this pre-Upper Cretaceous reflecting sequence in the Sirte Rise seems relatively thin.
- Both the seismic data and calculated magnetic basement indicate that in the Sirte trough the thickest and more complete sedimentary sequence of the Sirte Rise exists. The magnetic basement is expected here at about 7 to 8 km, including 1.8 km of water. From the Paleozoic to the Lower Jurassic a maximum total thickness of about 2.3 km has been calculated. Then, the Middle-Upper Jurassic to Oligocene sequence is generally thin (maximum 1.8 km in Sirte trough. The Miocene thickens progressively seaward and in the Ionian abyssal basin arrives at more than 3 km. The maximum Messinian thickness is about 1 km. The Plio-Quaternary is thin everywhere (100-400 m).
- The fault pattern of the Sirte Rise suggests that initial extensional movements commenced probably in the Middle Jurassic during the crustal opening of the Ionian Sea. But the main stretching phase and subsidence occurred in the Middle-Upper Cretaceous, when the prominent marine ingression of the Sirte basin took place. Also the "NQ" phase was active as indicates the fact that several faults affect the Tertiary Quaternary layers.
- From the Cyrenaica ridge to the Ionian abyssal basin, or from about 6.0 to 7.0 seconds on the top Mesozoic time structural map (Plate III), crustal conditions that can be attributed to an "intermediate type" exist. It is probably here that the SE continuation of the paleo-tectonic area, corresponding to the Sicily-Malta-Medina mounts escarpment, stretched during the "J" phase, and successively enormously modified by the younger extending processes, is to be found.

2.3 — Ionian Abyssal Basin

For several years, many investigations have been conducted in the Ionian Sea with the purpose of understanding the geological conditions and geodynamical evolution of such a key area of the Mediterranean.

In the past, when geophysical information was more scarce, the crustal conditions of the Ionian were strongly questioned. Successively, various Authors favoured the hypothesis that the Ionian basin (abyssal plain) is an old oceanic crust covered by a thick sedimentary sequence. But still continental crusts are supported by some Authors.

In the present scientific paper, the Author, after examination and interpretation of a great amount of geophysical geological and borehole data, summarizes his view.

Various data, interpretative elements, and arguments support the idea that the Ionian abyssal basin area is composed by a paleo-oceanic crust. Moreover, the interpretation of a regional seismic exploration covering the entire Mediterranean, and the numerous borehole data and seismic evidence on the occurred volcanic activities, indicate that crustal opening very probably took place during the Middle Jurassic.

To summarize, it is possible to list the following arguments in support of the existence of a paleo-oceanic crust in the Ionian abyssal basin:

- From the calibrated area of the Sirte basin and the Southern Pelagian Sea, seismic explorations allow some continuous clear reflectors to be followed up to the Ionian abyssal basin with high reliability regarding horizon identification. These reflectors indicate that the base of the Upper Cretaceous-Mesozoic sequence, which onlaps the basement and is fully calibrated in the Sirte basin, continues up to the Ionian abyssal basin with thinner sequences.
- On the Sirte slope, the Upper Cretaceous is underlain by older sediments having angular unconformity. These sediments thicken in the Sirte trough and again become relatively thin from the Cyrenaica ridge to the abyssal basin, along with the Lower Sirte slope.
- The acoustic basement is strongly fractured, not clear from the Sirte basin to the Cyrenaica ridge, and on the upper part of the area between the ridge and the abyssal basin. Then the basement becomes flat, more clear, and continuous except when recent volcanoes occur. This last basement seems more similar in character to that of an oceanic crust.
- From the calibrated area of SW Melita plateau, seismic reflectors, picked up to the Sirte rise and the Ionian abyssal basin, indicate that in the Sirte rise, under the Upper Cretaceous base (unconformity) older sediments attributable to a time-interval from Lower Cretaceous to Triassic exist. These sediments are thinning in the Sirte rise due to evident stretching and rifting processes which occurred during the "K" phase and probably also the "J" phase.
- The reflectors calibrated in the Sirte basin and in the Melita plateau area, and picked up to the abyssal basin, correlated across the Sicily-Malta escarpment, with the chronologically equivalent reflectors calibrated in the Sicily-Malta plateau area, indicate that, very probably, the lower part of the Mesozoic sequence is missing in the Ionian block.
- The studied area shows that the Middle Jurassic, everywhere on the Ragusa-Malta plateau, up to the Medina mounts, is affected by prominent volcanic activity constituted by widely diffused basaltic extrusions. Igneous rocks intervals are impressively continuous both in time and space on the greater part of the Middle Jurassic of the Sicily-Malta area, near the escarpment. Such activity is surely associated with a mega-geodynamic event that cannot be, logically, explained by a simple continental rifting process.
- The Bouguer gravity map shows that in the Ionian abyssal basin the absolute

IONIAN ABYSSAL BASIN




Fig. 14 — Example of Interpreted Seismic Reflection Line in Ionian Abyssal Basin (Line MS-27; S.P. 970-695). Extending the interpretation of Sirte Basin and Pelagian Sea Areas calibrated by boreholes to the Ionian Abyssal Basin, the hypothesis of Middle-Upper Jurassic sediments on a basaltic basement, fits very well on the entire seismic network of Ionian Basin. Evidence of extensional faults of "K" and "NQ" phases and of volcanic activity in Miocene.

IONIAN ABYSSAL BASIN




Fig. 15 — Example of Interpreted Seismic Reflection Line in Ionian Abyssal Basin (Line MS-27; S.P. 590-390). Evidence of faulting and volcanic activity associated with "K" and mainly "NQ" stretching phases. One volcanic show is outcropping with a small Sea Mount.

IONIAN ABYSSAL BASIN




Fig. 16 — Example of Interpreted Seismic Reflection Line in Ionian Abyssal Basin (Line MS-27; S.P. 230-1). Same conditions as previous example with volcanics in upper Miocene-Quaternary.




Fig. 17 — Example of Interpreted Seismic Reflection Line in Ionian Abyssal Basin (Line MS-21; S.P. 690-120). This part of Seismic Line MS-21 is located on the maximum Bouguer anomaly area (+ 310 mGal) of Central Ionian Basin. Evidence of prominent magmatic extrusions with a Sea Mount here named for the first time "Marconi Sea Mount" dedicated to one of the World's greatest scientists. Sedimentary sequence is considerably thinning and basement uplifting.




Fig. 18 — Example of Interpreted Seismic Reflection Line in Ionian Abyssal Basin Near the Outer Limit of Calabrian Arc (Line MS-60; S.P. 2020-2225). This and the next six Figures show the entire line MS-60, which crosses the whole Ionian part of Calabrian Arc. In this Figure are shown the typical flat conditions of the Ionian Abyssal Basin with sediments from Middle-Upper Jurassic to Quaternary.

maximum anomaly of the Mediterranean (Finetti-Morelli, 1973) of +310 mGal exists.

Such a very high value cannot be explained with models of continental crust or of intermediate types. Here, a very consistent thinning of the Mesozoic sequence and uplifting of the basement (*Fig. 17*) and prominent volcanic activities with a Sea mount — to which we give the name of "Marconi" — took place.

- h) Recent Deep Seismic Sounding investigations with seabottom seismographs indicate in the Ionian, off the maximum Bouguer anomaly, an oceanic crust with a Moho depth of about 17 km. This value is in agreement with our calculations and we expect a further crustal thinning at the Marconi sea mount area, up to a minimum of about 13-14 km (*Fig. 34*). This crustal model may now explain the Bouguer anomaly very well.
- i) Assuming a model which postulates a crustal opening in the Middle Jurassic as would indicate all the above listed arguments, all data used for this study fit very well, and in any case much better than models of continental or intermediate crusts.
- j) The Apulian platform, on its SW side, in the Ionian Sea, is affected by clear extensional geodynamics. The extended area with faults and back-tiltings is about 100 km wide. The latest movements occurred in recent times. Such a deformation pattern of the Apulian slope seems well in agreement with a model of a still moving passive margin.

Regarding the proposed crustal opening it is to be observed that the regional seismic exploration of the OGS here interpreted, does not indicate the existence of a mid-oceanic ridge, unless more detailed seismic control extends the Marconi feature north-westward and south-eastward. Probably, the hypothesis of the detachment and drifting of the Apulian (or Adriatic) plate, without mid-oceanic ridge creation, is the more correct interpretation.

2.4 — Apulian Platform

The Apulian platform is characterized by a very thick Mesozoic sedimentary sequence, continuous from the Permo-Triassic to the Upper Cretaceous in the southern part, and with the Upper Cretaceous missing in the northern one (about north of 41° parallel).

A deep borehole drilled in the Southern Apulia (Ugento 1) indicates about 4000 metres of Cretaceous over Jurassic. In the northern part (area of Canosa positive gravity anomaly) geophysical data indicate a very thin Lower Cretaceous (less than 200 metres) overlaying a Jurassic sequence about 4000 metres thick. So, in southern onshore Apulia we expect the top Triassic at about 8 km depth.

Lithologically, the Mesozoic succession is almost completely characterized by monotonous carbonate sequences, constituted by predominant limestone in the Upper Cretaceous and Dolomite in the Lower Cretaceous, Jurassic and Triassic. The Upper Triassic evaporite, well evident in the Adriatic from seismic data and constituted by halite in the Southern Adriatic basin and in the Peloponnesus area, on the contrary is not so evident in the Apulian seismic reflection data. From refraction seismics it is possible to discriminate in Apulia a very high velocity refractor of about 7.0 km/s, attributable to an anhydrite interval encountered in a borehole of the Gargano area at about 2200 metres below sea level, but much deeper in the Apulian peninsula.

Structurally, the emerged Apulian platform is rather tabular or very slightly undulated. The Gargano feature, very probably, is essentially a diapire-like structure generated by horizontal movements, with transform and shear faults along with the Triassic evaporitic interval. Clear seismic evidence of a transcurrent fault, crossing the Gargano and Southern Adriatic area exists for the Gargano fault (*Plate-III*). Evidence of

a regional transform fault can also be found for the Kefallinia fault which limits the Apulia platform and the Hellenic arc in the NE Ionian Sea.

The Apulian platform extends considerably south-eastward, up to the Kefallinia Island area. In the Adriatic it extends irregularly from about 20 to 45 km and then across a steep slope passes to much thinner sequences, probably corresponding to a deeper deposition after the "J" extensional phase of the Middle Jurassic. On the Ionian side clear evidence of extensional geodynamics exists (*Figs 4 & 5*). Here, the faulted slope connects the thick platform area with the relatively thin Mesozoic sequence of the Ionian abyssal basin interpreted as paleo-oceanic crust.

All the reconstructed structural data from the Apulian platform and surrounding vast area point to the convincing conclusion that Apulia, after its detachment from the Africa megaplate has drifted north-eastward and nowadays, due to the horizontal pressure of Dinarides and Hellenides on one side and of Apennine on the other, is moving anticlockwise along the Kefallinia and Gargano faults. Seismicity of the area is well in agreement and fully explained by such a postulated rotation.

2.5 — Southern Adriatic Basin

There is a consistent geological difference between the area of the central-northern Adriatic with shallow water and the area of the Southern Adriatic basin (*Plate III*) with deep water. A substantial analogy probably existed between these two areas from the Permo-Triassic until the Lower Jurassic, with a relatively thick dolomitic sequence and an evaporitic succession in the Upper Triassic. This evaporite is mostly anhydritic in the central northern Adriatic, while in the southern basin a thick halite precipitation is indicated by seismic data.

After the main "J" extensional phase of the Middle Jurassic, which occurred in the central-eastern Mediterranean, very probably, also the southern Adriatic, which is the marginal basin of the Apulian (Adriatic plate), was affected by the stretching activity with consequent subsidence and depositional change.

Geophysical data indicate that from the Jurassic (probably from the Middle-Upper Jurassic) until the end of the Oligocene the sedimentary sequence is relatively condensed. In the Miocene, due to the high deposition rate connected to the compressional phase of Dinarides, the sequence becomes progressively very thick from west to east (Finetti-Morelli, 1973: *Fig. 25*).

At the top of the Lower Cretaceous in the Southern Adriatic basin, as well as in the greater part of the Pelagian, Ionian and Eastern Mediterranean Seas, it is possible to identify a continuous characteristic good to fair reflector corresponding to the "Marne a Fucoidi" layer. This reflector is generally associated with a relatively thin interval of Marls interbedded with a carbonate sequence.

2.6 — Calabrian Arc

To observe the deformational process that took place in the Ionian part of the Calabrian Arc it is convenient to follow the entire interpreted Seismic line MS-60 from south-east to northwest (*Figs 18, 19, 20, 21, 22, 23 & 24*). The first figure (*Fig. 18*) shows the undeformed area of the Ionian abyssal basin just off the outer limit of the Calabrian arc. Here the sedimentary crust is the typical one, interpreted as paleo-oceanic, already seen in previous figures (*Figs 14, 15, 16 & 17*), and characterized by a relatively thin Mesozoic (about 1.3 sec reflection time), overlain by a thick tabular Tertiary Sequence (more than 2.4 sec) mainly constituted by Miocene.

Successive figures (*Figs 19 to 24*) show the arcuated area deformed by the Neogene-Quaternary compressional geodynamics. In *Fig. 19*, at the outer limit of the Calabrian arc, the shearing plane corresponds to the base of Messinian evaporite. This




Fig. 19 — *Example of Interpreted Seismic Reflection Line On the Most External Part of Calabrian Arc (Line MS-60; S.P. 1720-2000).*
 From flat conditions of Ionian Abyssal Basin one passes to the sliding interval, in front of the Arc and then to the beginning of the Fore Arc Scraping Zone.
 A shearing plane at the base of Messinian is evident and deepens progressively.

LINE MS 60

CALABRIAN ARC




Fig. 20 — *Example of Interpreted Seismic Reflection Line Across Calabrian Arc (Line MS-60; S.P. 1340-1620).*
 Fore Arc scraped blocks on a shearing plane progressively deepening inward, from about 6 to 8 seconds.

CALABRIAN ARC

LINE MS 60




Fig. 21 — Example of Interpreted Seismic Reflection Line Across Calabrian Arc (Line MS-60; S.P. 1000-1300).
Shearing plane limiting base of scraping disappears below the bottom of section, involving deeper layers of Mesozoic in deformation process.

CALABRIAN ARC

LINE MS 60




Fig. 22 — Example of Interpreted Seismic Reflection Line Across Calabrian Arc (Line MS-60; S.P. 680-975).
Evidence of imbrication of large blocks.

CALABRIAN ARC





Fig. 23 — Example of Interpreted Seismic Reflection Line Across Calabrian Arc (Line MS-60; S.P. 320-640). Probably Crystalline-Metamorphic and Mesozoic Blocks of European Plate are imbricated from S.P. 320 to about 630. Assuming such interpretation, S.P. 630 should correspond to the Trench Zone of Calabrian Arc, morphologically not so evident as the Hellenic Trench.

LINE MS 60

CALABRIAN ARC




Fig. 24 — Example of Interpreted Seismic Reflection Line Across the Calabrian Arc (Line MS-60; S.P. 20-300). Imbrication of large blocks which includes the Crystalline-Metamorphic and Mesozoic.

CALABRIAN ARC




Fig. 25 — Example of Interpreted Seismic Reflection Line Showing the Overthrusting of Calabria Arc on Ionian Abyssal Basin Area (Line MS-20; S.P. 2140-1610). On SW part of Calabrian Arc the slope of the underthrusting Ionian sequence is much more accentuated than on southern part (Compare e.g. with Figure 19).

plane then deepens progressively (Figs 20 & 21), involving the entire Tertiary sequence and the Upper Mesozoic. On Fig. 22 it is not possible to follow it any more because of the thick thrust blocks. Very probably, after about the S.P. 1000, as indicates also the Bouguer gravity gradient, the shearing plane enters the greater part of the total sedimentary sequence.

In the outer part of the arc (Fig. 19), from about S.P. 2000 to 1900, there is the frontal sliding allochthonous or melange. Such gravitative sliding deposits can be found in the foredeeps in front of every major thrust block, or where a certain slope of Messinian and soft sediments exists. From S.P. 1900 to about S.P. 630 (Figs 19 to 23) imbricated blocks detached from the underthrusting African plate are explored and shown. If our interpretation regarding the crustal opening of the Ionian is correct, we expect that these blocks generally do not include older sediments than Middle Jurassic, except possible continental fragments which persisted after the drifting of the Apulian plate.

So, the area from S.P. 1900 to 630 would correspond to the fore-arc scraping zone which seismically seems quite similar to other more classic scraping zones of the Indian and Pacific oceans. In the Calabrian arc the existence of a trench zone like for example, in the Hellenic arc is not immediately evident. But, a careful examination of the available geophysical data (seismic and gravity) combined with known geological, borehole and physiographic data may distinguish on Figs 23 & 24, from S.P. 630 to 20 the area where crystalline-metamorphic blocks and Mesozoic blocks of European plate are overthrust on the forearc scraping zone. In this case the trench zone of S.P. 630 is very close and associated with subhorizontal or slightly dipping fault planes. The sea bottom surface, across the interpreted trench zone, drops down about 0.4 sec. or about 300 m. (Plate 1).

While in the southern extremity of the Calabrian arc the Ionian crust is underthrusting with a very gently dipping plane (Figs 19 & 20), in the SW part (Fig. 25) as well as in the SE one, the Ionian sequence underthrusts the Calabrian arc with a much more accentuated slope. Observing these and other general aspects of the compressional deformation, it seems possible to argue that in recent geological time (Plio-Quaternary) the main component on the relative Africa-Europe movement in the Ionian Sea is not the N-S, but the SW-NE one. This fact determines now a shortening of the space of the Calabrian arc from Sicily-Malta escarpment to the Apulian margin with consequent restriction of arc width.

The same main component of the movement determines compression, overthrustings and shortening of the Ionian abyssal area in front of the Mediterranean ridge and of the ridge area. On the other side of the Hellenic arc, in the Levantine Sea, for this reason, the main consequent deformation is the gravity sliding.

2.7 — Hellenic Arc

The Ionian part of the Hellenic arc shows a regional structure very similar to that of the Calabrian arc. For a more complete geophysical documentation, in addition to the Figs 26, 27, 28, 29 & 30 we refer also to Finetti (1976).

In Fig. 26, Line MS-33 shows the structural and time stratigraphic relationships between the Ionian abyssal basin and the Mediterranean ridge. The Mediterranean ridge can be considered in summary the fore-arc scraping zone of the Hellenic arc, formed entirely by detached and imbricated blocks of sedimentary crust of the Ionian Sea. The Messinian, including the salt layer, is here relatively thick with respect to the Ionian Late Miocene evaporite sequence. In Fig. 28 the shearing plane coincides practically with the base of the Messinian salt. Then in (Fig. 29) entering more in the Mediterranean ridge, this plane deepens progressively and involves older sediments. Figs 27 & 30 show the structural conditions where relatively big blocks are imbricated.




Fig. 26 — Example of Interpreted Seismic Reflection Line Showing the Contact Between Ionian Abyssal Basin and Mediterranean Ridge (Line MS-33; S.P. 1680-1425).
Messinian interval "AB" is relatively thick and forms Salt Domes in front of the Scraped Zone. In this outermost part the shearing plane occurs along the Messinian interval.




Fig. 27 — Example of Interpreted Seismic Reflection Line Across Mediterranean Ridge (Line MS-33; S.P. 1440-1140).
It shows the typical imbrication with large blocks involved.




Fig. 28 — Example of Interpreted Seismic Reflection Line Showing the Contact Between the Cyrenaican Margin and the Mediterranean Ridge (Line MS-50; S.P. 12960-12660).
Thick Messinian (near one second) and beginning of Fore Arc Scraping Zone with shearing plane flat and sloping gently.




Fig. 29 — Example of Interpreted Seismic Reflection Line Across Mediterranean Ridge (Line MS-50; S.P. 12670-12400).
It shows the scraping process, and around S.P. 12470 the faulting commences to involve Mesozoic.




Fig. 30 — Example of Interpreted Seismic Reflection Line Across Mediterranean Ridge (Line MS-71; S.P. 460-670).
Typical compressional deformation with imbrication of blocks and deep faults.

In the Hellenic arc the existence of a trench zone (*Plate-III*) separating the thrust blocks of European plate over the scraped and imbricated blocks of the underthrusting African plate is well evident. From geophysical data it seems possible to say that the crustal layer, separating the imbricating blocks and the underthrusting, corresponds probably to the paleo-oceanic basement. In fact, no remarkable magnetic anomalies exist in the Mediterranean ridge. Therefore it seems that imbrication involves only sedimentary layers.

In *Plate II* the reconstructed geological cross-section of the Mediterranean ridge and the Ionian abyssal basin is shown. Moho discontinuity is calculated from gravity.

3. Volcanic activity

In the previous paragraphs, we have commented on the volcanic activities and their space distribution on the various studied geological provinces. For a more organic view, this important argument is now more specifically examined with the purpose of following the time and space distribution of volcanism and of understanding connected geodynamical meanings.

Assembling all the numerous known volcanic shows outcropping, checked by boreholes, or deduced from geophysical exploration, a function of igneous activity against geological time from Permo-Triassic until the present with an arbitrary scale is compiled (*Fig. 32*). In this figure the plotted intensity of volcanic activity is obtained taking into account the total thickness of igneous intervals occurring in a given province, in the considered geological time, and the space distribution density of the same area.

The following four main phases of igneous activity have been recognized in the Pelagian and Ionian Seas during about the last 230 M y.:

- 1) Middle-Upper Triassic ("T")
- 2) Middle-Jurassic ("J")
- 3) Middle-Upper Cretaceous ("K")
- 4) Middle-Upper Miocene to Quaternary ("NQ")

These phases are associated with corresponding paroxysmic periods of extensional geodynamics occurred on a more or less wide part of the studied area, or on a determined province of it. In *Fig. 31* the space distribution of the volcanic activities with indication of the corresponding extensional phase is shown.

This time-space distribution is based on all the most important data until now gathered. It is evident that further explorations may improve the knowledge. In some areas, scarcely known at present, it is possible to expect event consistent modifications, and/or additional activities with respect to the present information.

The diagram of *Fig. 32* indicates the four main extensional phases recognized. This does not mean that between a phase and the successive one completely quiet conditions occurred. A main phase is intended as a geological interval of time during which remarkable geodynamical (rifting) processes took place on a wide area or along a regional fault. In between each phase the geodynamic activity produced, in a given province, more moderate deformations, but the activity did not necessarily cease completely.

The first phase "T" produced its checked effects on the Sicily Malta area with frequent basaltic intervals interbedded with Middle-Upper Triassic sedimentary sequences (*Fig. 31 & Plate IV*). Very probably, also in the Gabes-Tripoli-Misurata basin a stretching activity occurred. We have no information, but we suppose activity of this phase also in the Sirte trough area.

As already mentioned above, in the Middle-Jurassic the most impressive and prominent volcanic activity of the Sicily-Malta area took place. Practically, on a large




Fig. 31 — Space-Time Distribution of Main Igneous Activities Checked by Boreholes and/or Inferred from Seismic and Magnetic Information.

MAIN EXTENSIONAL PHASES AND RELATIVE IGNEOUS ACTIVITY IN PELAGIAN AND IONIAN SEAS




Fig. 32 — Main Extensional Phases and Relative Igneous Activities of Pelagian and Ionian Seas.

part of this province, such a time stratigraphic interval is completely or almost completely composed by igneous rocks. In Malta Island along the Sicily-Malta escarpment Middle Jurassic volcanic rocks are found. Also on the deeper part of the Sirte rise this volcanic phase can be expected.

The existence of volcanic rocks of the Middle-Upper Cretaceous phase are proven in the following areas: Gabes-Tripoli-Misurata Basin; Sicily-Malta plateau area. Geophysical evidence of "K" phase volcanism can be clearly recognized in the Sirte rise and very probably in the eastern Medina bank, and in the Ionian abyssal basin too.

The last Neogene-Quaternary phase ("NQ") with maximum activity in Middle-Upper Miocene to Quaternary is the most widely diffused and generalized one. Proven "NQ" igneous rocks exist in Ragusa and Sicily-Malta area, and mainly in the Pantelleria-Linosa-Medina grabens area: basalts of Lentini; volcanoes of Pantelleria and Linosa; volcanics of the Medina Graben of Fig. 9. But, other numerous volcanic shows, all along with the rifting area of the Sicily Channel are evident from seismic and magnetic data. Clear and reliable seismic evidence of "NQ" volcanic activity is recognized in the Sirte rise, Medina Channel, Sicily-Malta escarpment. Impressive volcanic bodies are recognized in the Ionian abyssal basin (Figs 14, 15, 16 & 17). The most important one is that of the Marconi sea mount area (Fig. 17) where the thinnest crust conditions of the Ionian Sea exist.

4. Geodynamic evolution

In recent years, several plate tectonic models of evolution of the Mediterranean or part of it have been proposed by different Authors. Some of these are mentioned in paragraph 1. It is not the purpose of this paper to propose a new model or to analyze those already proposed and how they fit with the numerous regional data and conclusions here described. We wish only to observe that several basic propositions contained in some geodynamic models like, for example, in those of Biju-Duval et al., (1977), Laubscher et al., (1977), Scandone et al., (1976), and other Authors, are supported by relevant controlled data and arguments.

In the following, in greater detail, the most important geodynamical phases and corresponding deduced evolutive modifications of the investigated area, from the Permo-Triassic to the present, are commented on.

4.1 — Main extensional phases

As already mentioned in previous paragraphs, four main phases of geodynamical extensional activity, in the Central Mediterranean, from the Middle-Upper Triassic to the Quaternary, are recognized. These phases, obviously, correspond to those of the main volcanic activities:

- 1) Middle-Upper Triassic phase ("T")
- 2) Middle-Jurassic phase ("J")
- 3) Middle-Upper Cretaceous phase ("K")
- 4) Neogene-Quaternary phase ("NQ")

Analysis of extensional movements is limited to the northern part of the African plate and marginal basins.

4.1.1 — The Triassic continental rifting

Using in priority all numerous land or shallow water marine data available, utilizing also the relevant geophysical exploration of deep water on the Ionian Sea, it is possible to arrive at the fairly controlled conclusion that from the Late Permian to the Lower Triassic, sedimentation took place on the northern continental margin of Africa in a substantially quiet geodynamical condition.

In the Middle Triassic the first recognised extensional phase, which continued also during the entire Upper Triassic, commenced its activity. In particular three areas have been stretched and deformed by this "T" phase (Plate IV):

- a) Gabes-Tripoli-Misurata basin
- b) Streppenosa trough on Ragusa-Malta area
- c) Sicily-Malta-Medina mounts escarpment

The Gabes-Tripoli-Misurata basin already existed during the Paleozoic, when the Gafsa-Jeffara extensional fault system originated. During the Middle Triassic this basin was newly stretched with consequent subsidence and expansion.

From the southern part of the Ragusa platform to the Malta area, before the "T" phase, in the Lower Triassic, a generalized platform carbonate sedimentation took place. This original platform was stretched and rifted so as to produce a regional trough (Plate IV: Streppenosa trough), where sedimentation assumed the characteristics of restricted pelagic condition ("Streppenosa" Fmt., or Black Shale). Part of this pelagic sedimentation, was probably involved in the successive crustal opening of the Middle Jurassic, and drifted away. The Sicily-Malta-Medina mounts escarpment and the Lower Sirte slope, were also affected by the same extensional geodynamics. The numerous basaltic interval existing in the Ragusa-Malta area on the Middle-Upper Triassic support this hypothesis. Locally, movements continued also in the lower part of the Lower Jurassic, but, in general, diminished drastically or ceased almost completely at the end of the Triassic.

This "T" phase produced an important rifting process that, probably, stopped at the continental stage with wide seaways, thinnings, and subsidences. This opinion is favoured by several Authors: Biju-Duval et al., (1977); Scandone (1976); Hsü (1978), and others.

4.1.2 — The crustal opening of Ionian Sea and Eastern Mediterranean in Middle Jurassic

During the Middle Jurassic along with the Ionian-Eastern Mediterranean marging of the African plate, the most important extensional phase, testified by prominent volcanic activity, occurred in the Central-Eastern Mediterranean. More specifically, on the Sicily-Malta-Medina Mounts escarpment area and on the deeper part of the Sirte rise, the extensional phase "J" performed the complete opening of the Ionian Sea. It is impressive to note that this crustal opening is to the opening of the South Atlantic Ocean.

In the Ragusa-Malta plateau area, particularly on its eastern part, the Middle Jurassic interval consists almost completely or only of igneous rocks. The magnetic anomaly map indicates that this controlled "J" volcanic activity of the eastern Ragusa-Malta plateau, very likely, extends on the Medina mounts area and on the Lower Sirte rise.

During this phase, the Apulian platform together with the whole Adriatic plate, which were already separated from Africa by a wide sea channel, during the previous phase "T" moved consistently far away northeastward. The Sicily-Malta-Medina mounts steep escarpment separates the continental crust of the Ragusa-Malta plateau from the oceanic crust of the Ionian abyssal basin (Fig. 6). A scheme of crustal relationships between the two areas is given in Fig. 33.

Southeast of the Medina Mounts, the transition from continental crust to oceanic conditions, across the stretched, thinned, and subsided Sirte rise is much more progressive than across the Sicily-Malta escarpment. In fact, across this escarpment, at top Mesozoic (Plate III), we pass directly with a mega-fault system from 0.5 ÷ 1.4 seconds reflection time of the plateau area to the value of about 7.0 seconds of the abyssal basin in




Fig. 33 — Comparative Crustal Scheme of Pelagian Sea (Ragusa-Malta Plateau Area) and Ionian Basin.




Fig. 34 — Crustal Scheme of "Marconi Sea Mount" Area on the Maximum Mediterranean Gravity Anomaly. Basaltic basement remarkably uplifts and sedimentary sequence becomes much thinner. Prominent volcanic activity of "NQ" and very probably older phases.

only some 20 to 30 km. South of the Medina mounts the transition from shallow continental shelf to the Ionian abyssal basin takes place across the wide area of the entire Sirte rise, where completely different tectonic deformation occurred during the various geodynamical phases. East of the Medina plateau, for example, there is the Medina escarpment mostly due to the "K" and "NQ" stretching phases. This escarpment cannot be correlated with the Ragusa-Malta escarpment because of the enormously different dimensions and geological conditions. If we want to look for the southeast continuation of the equivalent tectonic feature of the Ragusa-Malta-Medina mounts escarpment, we can find it in the area from the Lower Sirte slope to about the 7.0 seconds reflection time (*Plate III*) at top Mesozoic. This area, now characterized by an intermediate crust type, very close to ocean like conditions, was greatly stretched during the "J" phase and, probably also during the "T" one. Successive "K" and "NQ" phases contributed to stretch the entire Sirte rise area and again the lower slope and deeper area above mentioned.

Therefore, any tentative correlation between the Ragusa-Malta escarpment and the Medina escarpment has very scarce geological meaning, because there is a much more significant correlation between the Ragusa-Malta-Medina mounts escarpment and the lower Sirte slope area (*Plate III*). The Medina mounts, after the various geodynamical deformations that passed through show now a complex extensional tectonics with several fragmented blocks and collapsing faults having two main trends: E-W and NE-SW. Some shear faults seem to occur too.

4.1.3 — Stretching Cretaceous phase and associated prominent transgression

From the end of the Middle Jurassic to the Lower Cretaceous included, no remarkable and diffused volcanic activity or crustal movements seem to have affected the studied area, except for an evident subsidence in the Early Cretaceous (Neocomian transgression).

At the end of the Lower Cretaceous, a relatively thin, widely diffused marly interval ("Marne a Fucoidi" layer) all over the studied Central Mediterranean area, interbedded in carbonate sequence, occurs. This layer is characterized by a good and continuous seismic reflector and is probably associated with a subsidence or with a sea level uplift. Extensional geodynamic movements with subsidence and a remarkable transgression characterize the Upper Cretaceous (Cenomanian transgression) which is very prominent in the Northern Africa area, but is also evident in the Apulian and Adriatic areas. Thus, the "K" phase of our scheme of *Fig. 32* produced stretching, faulting or activation of existing faults, subsidence, and magmatic activities of a basaltic or pyroclastic type in several parts of the Central Mediterranean.

In the Gabes-Tripoli-Misurata basin there is clear evidence of extensional tectonics with faulting and basaltic effusions widely distributed in the Late Cretaceous. Volcanic activity of the "K" phase is also found in the Ragusa-Malta plateaus area and more sporadically in other parts of the Pelagian Sea. Evidence of tectonic and volcanic activity of this phase exists also in the Ionian abyssal basin.

But, the geological province where the "K" phase seems to have produced its most remarkable effects is that of the Sirte rise and the Sirte basin. In the Sirte rise the stretching activity is proved by the existence of numerous faults with back-tilted blocks and volcanic effusions of clear Upper Cretaceous age. This geodynamics produced a consistent subsidence of the whole Sirte rise and of the Sirte basin, and a remarkable marine ingression on the north African plate. Apulia continued its moving away.

4.1.4 — The Miocene-Quaternary phase and the huge rifting of Sicily Channel

The last extensional phase was active from the Middle Upper Miocene to the Quaternary. It is possible to recognize volcanic extrusions in many zones of the Pelagian and Ionian Seas. Well known volcanoes are those of Pantelleria and Linosa Islands. Volcanic outcrops associated to this phase exist in the Ragusa area (Basalts of Lentini, etc.).

The seismic evidence of faulting and of igneous bodies of the Neogene-Quaternary age is identified on several sites of the Pantelleria-Linosa-Malta-Medina grabens area. See *Fig. 9* for example. Other shows are found in the Sicily-Malta escarpment (*Fig. 6*) and south of the Medina mounts (*Fig. 7*).

Very clear seismic evidence of "NQ" volcanic activity and faulting exists in numerous points of the Ionian abyssal basin (*Figs 14, 15, 16 & 17*) and of the Sirte rise.

All these data show that the "NQ" phase produced its effects on a vast part of the studied area. However the area where this phase determined the most remarkable geological, geomorphological and even crustal modifications is that of the Pantelleria-Linosa-Malta-Medina grabens, or the area of the impressive rifting of the central Pelagian Sea, called globally "Sicily Channel" (*Fig. 1*).

The rifting process of the Sicily Channel is more developed in the area from Pantelleria-Linosa to Malta (*Figs 11, 12 & 13*) than at the NW extremity of the Pelagian Sea between Cape Bon in Tunisia and Mazara del Vallo in SW Sicily. Also along with the Medina graben the process is not so impressive as that shown on the seismic line MS-19. But the Sicily channel rifting area is continuous all along the entire Pelagian Sea from the Tunisian extremity to the Ionian Sea.

Seismic, gravity and magnetic data show clearly that the rifting, still active, has already produced a remarkable geological deformation involving not only the entire sedimentary sequence, but remarkable tilting movements on both sides of the Sicily channel area show that also the Lower crust participates in the geodynamical processes with uplifting of the earth's mantle. Bouguer gravity (*Fig. 2*) confirms this observation. In fact along the Sicily Channel there is a remarkable positive regional anomaly which, probably, is associated with a crustal thinning produced by the "NQ" stretching phase. It can be said that the rifting process has already evolved to such a stage as to practically now divide the Pelagian Sea essentially into two separated blocks. One to the north is formed by the Adventure and Ragusa-Malta plateaus and the second one on the south is formed by the Lampedusa and Medina plateaus. This second block remains substantially connected to the North-African plate, because even though it is affected by several extensional faults and grabens, these are much less important than those of the Sicily Channel.

The Apulian (Adriatic) plate has continued to move more moderately away from the African one and during this period its SE passive margin, in the Ionian Sea, has been affected by numerous new extensional faults or activation of previously existing ones. This extensional process is still active. A very high seismicity in recent and/or historical time has been observed along with the Kefallina and the Gargano regional transcurrent faults. Examining these data and the deformation pattern of the whole area surrounding the Apulian plate (Apennine, Dinarides, and Hellenides) it seems possible to argue that the Apulia is now moving anticlockwise along with the above mentioned Kefallina and Gargano faults.

4.2 — Compressional phase of Calabrian and Hellenic Arcs

In the Lower-Middle Miocene, the North African Margin commenced to collide with

the southern margin of the European plate. In particular the north Sicily margin disappeared beneath the European south-moving overthrusts.

At the same time the inner Calabrian arc composed by European blocks moved outward (SE) and overthrust the Ionian abyssal basin Crust. The Taormina transform fault has played an important role in this process.

The final result is a tectonically complex orogenic arc with overthrusts, reverse faults, repeated imbrication of blocks and gravity sliding (Fig. 19). In the Ionian the sediments covering the paleo-oceanic Crust of the African margin are scraped with a shearing plane that progressively deepens towards the inner arc.

According to our interpretation, the accretionary zone, out of the European thrust one, is composed by sediments from the Middle-Upper Jurassic to the Quaternary.

The Hellenic arc shows a deformational scheme similar to that of Calabria. On its Ionian side the compressional movements and the relative deformations are much clear than on the Levantine one.

As already observed in the Calabrian arc, also in the Hellenic one, the first deformational process, at the limit with the undeformed Ionian abyssal basin, is due to an imbrication of blocks of Messinian to Plio-Quaternary rocks over a shearing plane which corresponds to the base of the Messinian salt (Fig. 26 & 28). Then the overthrusting progressively involves older sediments and the shearing plane deepens (Figs 27, 29 & 30).

From seismic data a regional geological section across the Ionian abyssal basin and the whole Mediterranean ridge has been reconstructed (Plate II). The feature of the Mediterranean ridge, substantially, constitutes the fore-arc scraping zone of the Hellenic arc. It is formed by imbrication of blocks which include sediments sheared from the paleo-oceanic basement of the subducting African plate. If the deduced time of crustal opening of the Ionian (and the Eastern Mediterranean) is correct, here too, as in the outer Calabrian arc, the oldest sediments involved in the thrusting process would be those of the Middle-Upper Jurassic, but the existence of fragments of the opened African continental crust, including the Permo-Triassic and the Liassic, cannot be completely excluded.

The seismic stratigraphy clearly indicates that between a block and the previous or the successive one of the scraping zone of the Mediterranean ridge there is generally a possible correlation, while the Hellenic trench separates two zones completely different from the viewpoint of the seismic characters. This is perfectly comprehensible if it is assumed that the Hellenic trench is the regional tectonic contact between the subducting African plate and the European one.

Acknowledgments. The Author would like to thank the entire staff of the OGS's Seismic Division for the enormous and invaluable work carried out over several years of systematic exploration of the whole Mediterranean.

Many thanks are due to G. Bidussi, D. Calligaris, G. Cova, A. Diqua, G. Dordolo, G. Moimas, G. Renner, A. Scotti, G. Vascotto, L. Zampieri, V. Brandolin, R. Briseik, C. Coderin, R. Furlan, G. Morgan, G. Perini, L. Rocco, A. Salemm, and E. Tercovich for their contribution to the field seismic data acquisition.

Particular thanks are due to Prof. R. Nicolich, Ing. S. Sancin, Ing. S. Marchetti, Ing. L. Salvador, M. Marchi, F. Ponton, and R. Sinceri for the seismic data processing.

Thanks are also due to G. Cantar and M. Pisani of the OGS for the important gravimetric and magnetic data produced on the whole Mediterranean, and to Dr. M. Piccolo, Dr. V. Zucconi Galli Fonseca, and R. Polla Zappador of the Istituto di Geodesia e Geofisica of the University of Trieste for their invaluable help in the preparation of the various geophysical illustrations used in this paper.

The present paper is the publication n. 201 of the Istituto di Geodesia e Geofisica dell'Università di Trieste; it was presented at the XXVII Congress of I.C.S.E.M., Plenary Assembly, October 9-18, 1980, Cagliari, Italy and at the International Symposium on the Hellenic Arc and Trench, April 8-10, 1981, Athens, Greece.

Bibliography

- Aadland A.J., Hassan A.A., 1972. *Hydrocarbon potential of the Abu Gharadiq basin in the Western Desert*. Arab Republik of Egypt. Eighth Arab Petrol Congr., Algiers. Paper no. 81 (B-3).
- Abbate E., Bortoletti V., Passerini P., Sagri M., Sestini G., 1970. *Development of the northern Apennines Geosyncline*. Sediment. Geol., v. 4, p. 3-4.
- Adamia S.A., 1975. *Plate tectonics and the evolution of the Alpine system: Discussion*. Geol. Soc. Amer. Bull., v. 86, p. 719-720.
- Adams C.G., Ager D.V., eds., 1967. *Aspects of Tethyan Biogeography*. London: The Systematics Association.
- Afanassiev G.D., 1960. *A few traits on the Magmatism of Egypt*. Izv. Akad. Nauk. USSR, Ser. Geol. n. 3, p. 26-40.
- Akal T., 1972. *The general geophysics and geology of the Strait of Sicily*, in: Oceanography of the Strait of Sicily. Allan T.D., Akal T., Molcard R., eds. Saclant ASW Res. Cent., La Spezia, Conf. Proc. v. 7, p. 177-192.
- Alinat J., Hinz K., Leenhardt O., 1970. *Quelques profils en sondage sismique continuen Méditerranée occidentale*. Rev. Inst. Franç. Pétrole, v. 25, p. 305-326.
- Alla G., Dessolin D., Leenhardt O., Pierrot S., 1972. *Données du sondage sismique continu concernant la sédimentation Plio-Quaternaire en Méditerranée nord-occidentale*. In: The Mediterranean Sea — A natural Sedimentation Laboratory. Stanley D.J., ed., Stroudsburg: Dowden, Hutchinson and Ros, p. 471-487.
- Alla G., Leenhardt O., 1971. *Découverte d'un dome de sel (Dome T, Sud Iculon), avec le bathyscaphe*. C.R. Acad. Sci. Paris, v. 272 (D).
- Allan T.D., Akal T., Molcard R., eds., 1972. *Oceanography of the Strait of Sicily*. Saclant ASW Res. Cent., La Spezia, Conf. Proc. 7, 227 pp.
- Allan T.D., Morelli C., 1971. *A geophysical study of the Mediterranean*. Boll. Geof. Teor. Appl., v. 13, p. 99-142.
- Allan T.D., Pisani M., 1970. *Gravity, magnetic and depth measurements in the Ligurian Sea*. Rapp. C.I.E.S.M., v. 18, p. 907-909.
- Altinli E., 1963a. *Explanatory text of the geological map of Turkey, sheet Cizre*. Bull. Miner. Res. Explor. Inst. Turk., p. 51-103.
- Altinli E., 1963b. *Explanatory text of the geological map of Turkey, sheet Erzurum*. Bull. Miner. Res. Explor. Inst. Turk., p. 55-103.
- Altinli E., 1964. *Explanatory text of the geological map of Turkey, sheet Van*. Bull. Miner. Res. Explor. Inst. Turk., p. 41-90.
- Alvarez W., 1972. *Rotation of the Corsica-Sardinia microplate*. Nature Phys. Sci., v. 235, p. 103-105.
- Alvarez W., 1973. *The application of plate tectonics to the Mediterranean region*. In: Implications of Continental Drifts to the Earth Sciences. Tarling D.S., Runcorn S.K., eds., New York, Academic Press, p. 893-308.
- Alvarez W., 1974. *Sardinia and Corsica, one microplate or two?* Rend. Sem. Fac. Sci. Univ. Cagliari, p. 1-4.
- Alvarez W., Cocozza T., Wezel F.C., 1974. *Fragmentation on the Alpine orogenic belt by microplate dispersal*. Nature, v. 248, p. 309-314.
- Alvarez W., Franks S.G., Nairn J.E.M., 1973. *Paleomagnetism of Plio-Pleistocene basalts from Northwest Sardinia*. Nature Phys. Sci., v. 243 (123), p. 10-11.
- Alvarez W., Gohrbandt K.H.A., 1970. *Geology and History of Sicily*. Tripoli: Petrol Expl. Soc. Libya.
- Amin M.S., 1961. *Subsurface features and oil prospects of the Western Desert*. Egypt, U.A.R., Third Arab Petrol. Congr., Alexandria.
- Amodio-Morelli M., Bonardi G., Colonna V., Dietrich D., Giunto G., Ippolito F., Liguori V., Lorenzoni S., Paglionico A., Perrone V., Piccarretto V., Russo M., Scandone P., Zanettin-Lorenzoni E., Zuppetta A., 1979. *L'arco calabro-peloritano nell'orogene appenninico-maghrebide*. Mem. Soc. Geol. Ital., v. 17, p. 1-60.
- Andrawis S.F., 1972. *New biostratigraphic contribution for the upper part of the Paleozoic rocks of Gibb Afia Well No. 2, Western Desert, Egypt*. Eighth Arab Petro. Congr., Algiers, Paper n. 76 (B-3).
- Andrieux J., Fonthote J.M., Mattauer M., 1971. *Sur un modèle explicatif de l'arc de Gibraltar*. Earth. Planet. Sci. Lett., v. 12, p. 191-198.
- Angelier J., 1975. *Sur le plateformés marines quaternaires et leurs déformations: les rivages méridionaux de la Crête orientale (Grèce)*. C.R.Acad. Sci. Paris, v. 281 (D), p. 1149-1152.
- Angelier J., 1976. *Sur la néotectonique de l'arc égéen externe: les failles normales est-ouest et l'extension subméridienne crétoise*. C.R. Acad. Sci. Paris., v. 282 (D), p. 413-415.

- Angelier J., 1978. Tectonic evolution of the Hellenic Arc since the late Miocene. *Tectonophysics*, v. 49, p. 23-46.
- Angelier J., Lepichon, X., 1980. La subduction hellénique et l'expansion égeenne: reconstitution cinématique et interprétation dynamique. *Compte Rendu Sommaire des Séances, Bulletin de la Société géologique de France*, 22, (5), 158-161.
- Arana V., Vegas R., 1973. Plate tectonics and volcanism in the Gibraltarc Arc. *Tectonophysics*, v. 24, p. 197-212.
- Argyriadis I., Brunn J.H., Ricou L.E., de Graciansky P.C., Poisson A., Marcoux J., 1975. *Elements majeur de liaison entre Taurides et Hellénides*. (Abstract). Orsay, France: Vème Colloque sur la Géologie des Régions Egéennes.
- Arthaud F., Mattauer M., 1972. Présentation d'une hypothèse sur la genèse de la virgation pyrénéenne du Languedoc et sur la structure profonde du Golfe du Lion. *C.R. Acad. Sci. Paris*, v. 274 (D), p. 524-527.
- Aruta L., Buccheri, G., 1976. Biostratigraphy and paleoecology of the early Messinian carbonates (Baucina FM) in western Sicily. *Mem. Soc. Geol. IT.*, v. 16, p. 141-152.
- Assereto R., Bosellini A., Fantini Sestini N., Sweet W.C., 1973. The Permian-Triassic boundary in the Southern Alps (Italy), in Permian and Triassic Systems and Their Mutual Boundary. Logan A., Hills L.V., eds. *Calgary: Can. Soc. Petr. Geol.*, p. 176-199.
- Aubouin J., 1960. Essai sur l'ensemble italo-Dinarique et ses rapports avec l'arc alpin. *Bull. Soc. Géol. France*, v. 2, p. 487-526.
- Aubouin J., 1963b. Esquisse paléogéographique et structurale des chaînes alpines de la Méditerranée moyenne. *Geol. Rundsch.*, v. 53, p. 480-534.
- Aubouin J., 1970. Réflexion sur la tectonique de faille plio-quaternaire. *Geol. Rundsch.* v. 60, p. 833-848.
- Aubouin J., 1972. Chaînes liminaires (Andines) et chaînes géosynclinales (Alpines). 24th In. *Geol. Congr.*, Sect. 3, p. 438-461.
- Aubouin J., 1973. Paléotectonique, tectonique, tarditectonique et néotectonique en Méditerranée moyenne: à la recherche d'un guide pour la comparaison des données de la géophysique et de la géologie. *C.R. Acad. Sci. Paris*, v. 27 (D), p. 457-460.
- Aubouin J., Bosellini A., Cousin M., 1965. Sur la paléogéographie de la Vénétie au Jurassique. *Mem. Geol. Univ. Ferrara*, v. 1, fasc. II, p. 147-158.
- Aubouin J., Brunn J.H., Celet P., Decourt J., Geodfriaux I., Mercier J., 1963. Esquisse de la géologie de la Grèce. *Livre jubilaire P. Fallot. Mém. Soc. Géol. France, Hors-ser.*, v. 2, p. 583-10.
- Aubouin J., Dercourt J., 1970. Sur la géologie de l'Egée: Regard sur le Dodécanèse méridional (Kasos, Karpathos, Rhodes). *Bull. Soc. Géol.*, p. 453-572.
- Aubouin J., Dercourt J., 1975. Les transversales dinariques dérivent-elles de paléofailles transformantes? *C.R. Acad. Paris, Sér. D.*, p. 347-350.
- Auzende J.M., 1971. La marge continentale tunisienne: résultats d'une étude par sismique réflexion: sa place dans le cadre tectonique de la Méditerranée occidentale. *Mar. Geophys. Res.*, v. 1, p. 162-177.
- Auzende J.M., Olivet J.L., Bonnin J., 1972b. Une structure compressive au nord de l'Algérie? *Deep-Sea Res.*, v. 19, p. 148-155.
- Auzende J.M., Olivet J.L., Pautot G., 1973b. Structural framework of selected regions of the western Mediterranean: Balear Islands, southern prolongation. In: *Initial Reports of Deep Sea Drilling Project*, v. 13, p. 1441-1447.
- Auzende J.M., Olivet J.L., 1974. Structure of the Western Mediterranean Basin. In: *The Geology of Continental Margins*. Burk C.A., Drake C.L., eds., New York: Springer-Verlag, p. 723-731.
- Auzende J.M., Rehault J.P., et al., 1975. Le bassins sédimentaires de la mer d'Alboran. *Bull. Soc. Géol. France*, v. 17, p. 98-107.
- Awad G.H., Fawzi I.M.A., 1956. The Cenomanian transgression over Egypt. *Bull. Inst. Désert Egypt*, v. (1), p. 169-184.
- Avedik F., Hieke W., 1981. Reflection seismic profiles from the central Ionian Sea (Mediterranean) and their geodynamic interpretation. "Meteor" *Forsch. Ergebnisse Reihe*, No. 34, Seite 49-64.
- Badrawi R., 1963. The geological significance of the regional gravity data, eastern Mediterranean region. *Proc. Fourth Arab Petrol. Cong.*, Beirut, Paper 25 (B2), p. 6.
- Baird D.W., 1967. The permo-carboniferous of Southern Tunisia. In: *Guidebook to the Geology and History of Tunisia*. Petrol. Expl. Soc. Libya, 9th Ann. Field. Conf., p. 85-104.
- Baird D.W., 1970. Review of Mediterranean Alpine tectonics. In: *Geology and History of Turkey*. Campbell A.G., ed., Petrol. Expl. Soc. Libya, 13th Ann. Field. Conf., p. 139-158.
- Baldacci F., Elter P., Giannini E., Giglia G., Lazzarotto A., Nardi R., Tongiorgi M., 1967. Nuove osservazioni sul problema della falda toscana e sulla interpretazione dei flysch arenacei tipo "Macigno" dell'Appennino Settentrionale. *Mem. Soc. Geol. IT.*, v. 6, p. 213-244.
- Ballard R.D., Uchupi E., 1975. Triassic rift structure in Gulf of Maine. *Amer. Assoc. Petr. Geol. Bull.*, v. 59, p. 1041-1072.

- Barakat M.G., 1970. A stratigraphical review of the Jurassic formations in Egypt and their oil potentialities. Seventh Arab Petrol. Congr., Kuwait, Paper no. 58 (B-3).
- Barakat M.G., Aboul Ela N.M., 1970. Microfacies and paleoecology of Middle Eocene and younger sediments in Geneifa area, Cairo-Suez district. *J. Geol. U.A.R.*, v. 14 (1), p. 1-51.
- Barbieri F., Civetta L., Gasparini P., Innocenti P., Scandone R., Villari L., 1974. Evolution of a section of the Africa-Europe plate boundary paleomagnetic and volcanological evidence from Sicily. *Earth Planet. Sci. Lett.* p. 123-132.
- Barbieri F., Gasparini P., Innocenti P., Villari L., 1973. Volcanism of the Southern Tyrrhenian Sea and its geodynamic implications. *J. Geophys. Res.*, v. 78, p. 5221-5232.
- Baroz F., Benoulli D., Biju-Duval B., Bizon G., Bizon J.S., Letovzey J., 1978. Correlations of the neogene formations of the Florence rise and Northern Cyprus: Paleogeographic and structural implications. *Initial Reports of the Deep Sea Drilling Project*, v. 42, Part. 1, p. 903-921.
- Bartolini C., Malesani P.G., Manetti P., Wezel F.C., 1975. Sedimentology and petrology of Quaternary sediments from the Hellenic Trench, Mediterranean Ridge and the Nile Cone from D.S.D.P., Leg. 13, Cores., *Sedimentology*, N. 22, p. 205-236.
- Basta E.Z., Zaki M., 1961. Geology and mineralization of Wadi Sikeit area, south Eastern Desert. *J. Geol. U.A.R.*, v. 5 (1), p. 1-37.
- Bebien J., Blanchet R., Cadet J.P., Charvet J., Chorowicz J., Lapierre H., Rampnoux J.P., 1977. The volcanisme triassique des Dinarides. Sa place dans l'évolution géotectonique mésogéenne. *Tectonophysics*.
- Bellini E., 1969. Biostratigraphy of the "Al Jaghbub (Giarabub) Formation in Eastern Cyrenaica, Libya. *Proc. Third African Micropaleontol. Colloq.*, Cairo, p. 165-183.
- Bernoulli D., Jenkyns H.C., 1974. Alpine, Mediterranean, and Central Atlantic Mesozoic facies in relation to the early evolution of the Tethys. In: *Modern and Ancient Geosynclinal Sedimentation*. Dott R.H., Jr. Shaver R.H., eds., *Econ. Paleont. Mineral. Spec. Publ.* 19, p. 129-160.
- Bernoulli D., Laubscher H., 1972. The palinspastic problem of the Hellenides. *Eclogae Geol. Helv.*, v. 65, p. 107-118.
- Beydoun Z.R., 1972. A new evaluation of the petroleum prospects of Lebanon with special reference to the pre-Jurassic. *Proc. Eight Arab. Petrol. Congr.*, Algiers, Paper 80 (B-3), 15 p.
- Beydoun Z.R., Dunnington H.V., 1975. *The Petroleum Geology and Resources of the Middle East*. England: Beaconsfield, Scientific Press 99 p.
- Biely A., Burolet P.F., Ladjmi T., 1972. Etude géodynamique de la Tunisie et des secteurs voisins de la Méditerranée, 23e Congrès C.I.E.S.M., Athènes, 3-11 Novembre 1972, 9 pp.
- Biju-Duval B., 1974. Carte géologique et structurale des bassins tertiaires du domaine méditerranéen: commentaires. *Rev. Inst. Franç. Pétrole*, v. 28, p. 607-639.
- Biju-Duval B., Dercourt J., Le Pichon X., 1977. From the Tethys Ocean to the Mediterranean Seas: A plate tectonic model of the evolution of the western Alpine system. In: *Biju-Duval B., Montadert L. (eds.), Symposium on the geological history of the Mediterranean basins: Paris (Technip)*, p. 143-164.
- Biju-Duval B., Letouzey J., Montadert L., et al., 1974. Geology of the Mediterranean Sea basins. In: *The Geology of Continental Margins*. Burk C.A., Drake C.L., eds., New York: Springer-Verlag, p. 695-721.
- Biju-Duval B., Letouzey J., Montadert L., Courrier p., Mugniot J.F., Sancho J., 1974. Geology of the Mediterranean sea basins. In: *The Geology of Continental Margins*. Burk C.A., Drake C.L., eds., New York: Springer-Verlag, p. 695-721.
- Biju-Duval B., Letouzey J., Montadert L., 1978. Structure and evolution of the Mediterranean basins. *Initial Reports of the Deep Sea Drilling Project*, v. 42, Part 1, p. 951-981.
- Biju-Duval B., Montadert L., 1977. Introduction to the Structural History of the Mediterranean Basins. In: *International Symposium of the Structural History of the Mediterranean Basins, Split, Yugo.*, p. 1-12.
- Bishop W.F., 1975. Geology of Tunisia and adjacent parts of Algeria and Libya. *Amer. Assoc. Petr. Geol. Bull.*, v. 59, p. 413-450.
- Blanc-Vernet L., Chamley H., Froget C., Le Boulicaut D., Monaco A., Robert C., 1975. Observations sur la sédimentation marine récente dans la région siculo-tunisienne. *Géol. Médit.*, v. 2 (1), p. 31-48.
- Bobier C., Martin G., 1977. Remarques sur l'évolution cénozoïque du détroit siculo-tunisien. *Rapport. Comm. Int. Mer Médit.*, 24, 7a, p. 223-225.
- Boccaletti M., 1975. Plate tectonics model for the evolution of the western Mediterranean. *Geol. Balcanica*, v. 5, p. 19-28.
- Boccaletti M., Elter P., Guazzone G., 1971. Plate tectonic models for the development of the western Alps and northern Apennines. *Nature Phys. Sci.*, v. 49, p. 108-111.
- Boccaletti M., Guazzone G., 1974. Remnant arc and marginal basins in the Cainozoic development of the Mediterranean. *Nature*, v. 252, p. 18-21.
- Boer J., de, 1965. Palaeomagnetic indications of megatectonic movements in the Tethys. *J. Geophys. Res.*, v. 70, p. 931-944.

- Bonnefous J., 1967. *Jurassic stratigraphy of Tunisia: a tentative synthesis (northern and central Tunisia, Sahel and Chotts areas)*. In: Guidebook to the Geology and History of Tunisia. Martin L., ed., Tripoli: Petr. Explor. Soc. Libya, p. 109-130.
- Borsetti A.M., Colantoni P., 1973. *Stratigrafia dell'isola di Lampedusa (canale di Sicilia)*. Giornale di Geologia, (2) 39, fasc. 1, p. 313-320, t. XXX, Bologna.
- Borsetti A.M., Colantoni P., 1975. *Cretaceous and Eocene sediment dredged in the strait of Sicily*. Lab. Geol. Marina, Bologna.
- Bosellini A., 1973. *Modello geodinamico e paleotettonico delle Alpi Meridionali durante il Giurassico-Cretacico. Sue possibili applicazioni agli Appennini*. In: Moderne Vedute sulla Geologia dell'Appennino. Roma: Accad. naz. Lincei, p. 163-213.
- Bosellini A., Winterer E.L., 1975. *Pelagic limestone and radiolarite of the Tethyan Mesozoic: A genetic model*. Geology, v. 3, p. 279-282.
- Bowen C.F., Vondra C.F., 1974. *Paleoenvironmental interpretations of the Oligocene Gabal El Qatrani Formation, Fayum Depression, Egypt*. Ann. Geol. Surv., Cairo, v. 4, p. 115-137.
- Brinkmann R., 1974. *Geologic relations between Black Sea and Anatolia*. In: The black Sea Geology, Chemistry and Biology. Degens E.T., Ross D.A., eds., Amer. Assoc. Petrol. Geol. Mem. 20 p. 63-76.
- Burrollet P.F., 1964. *Remarques sur le Néogène de Tunisie et de Libye et ses relations avec les bassins voisins*. Instituto "Lucas Mallada", C.S.I. Espana, Cursillon y Conferencias, IX, pp. 199-202.
- Burrollet P.F., 1967. *General geology of Tunisia*. In: Guidebook to the Geology and History of Tunisia. Petrol., Expl. Soc. Libya, 9th Ann. Field Conf., p. 51-58.
- Burrollet P.F., 1981. *Structure and petroleum potential of the Ionian Sea*. Deep Offshore Technology Conference Proceedings. 19/22 October 1981. Palma de Mallorca, Spain. N. 1, p. 1-11.
- Burrollet P.F., Byramjee R.S., 1974. *Reflexions sur la Tectonique Globale, Exemples Africains et Méditerranéens*. Notes et Memoires Comp. Française Petrole, 11, p. 71-120.
- Burrollet P.F., Byramjee R., 1974. *Evolution of géodynamique néogène de la Méditerranée occidentale*. C.R.Acad. Sci., Paris, v. 278 (D), p. 1321-1324.
- Burrollet P.F., Clairefond P., Winnock E. (eds), 1979. *Geologie Méditerranéenne: La mer pelagienne*. Annal de l'Université de Provence. Tome VI, No. 1. Marseille.
- Burrollet P.F., Dufaure P., 1972. *The Neogene series drilled by the Mistral No. 1 well in the Gulf of Lion*. In: The Mediterranean Sea — A Natural Sedimentation Laboratory. Stanley D.J., ed., Stroudsburg Dowden, Hutchinson, Ross, p. 91-98.
- Burrollet P.F., Manderscheid G., 1965. *Le Crétacé inférieur en Tunisie et en Libye*. Colloque sur le Crétacé inférieur. Mem. B.R.G.M. no. 34, pp. 785-794, Paris.
- Caire A., 1970. *Sicily in its Mediterranean setting, in Geology and History of Sicily*. Alvarez W. Gohrbandt. K. and Gohrbandt K. (eds.). Petroleum Exploration Society of Libya, Tripoli p. 145-170.
- Caire A., 1978. *The Central Mediterranean Mountain Chains in the Alpine Orogenic Environment, the Ocean Basins and Margins*. Western Mediterranean, v. 4B, 1.
- Caputo M., Panza G.F., Postpischl D., 1970. *Deep structure of the Mediterranean Basins*. J. Geophys. Res., v. 75, p. 4919-4923.
- Carey S.W., 1976. *The expanding earth*. In: Development in Geotectonics, 10, 1-488, Elsevier Publ. Co., Amsterdam.
- Castellarin A., 1972. *Evoluzione paleotettonica sinsedimentaria del limite tra "piattaforma veneta" e "bacino lombardo" a nord di Riva del Garda*. Giorn. Geol. v. (2) 38, p. 11-212.
- Castellarin A., 1977. *Ipotesi paleogeografica sul bacino del Flysch Sudalpino Cretacico*. Boll. Soc., Geol. Ital., 95 (1976): 501-511.
- Castellarin A. 1980. *Jurassique et Crétacé*. Introduction à la Geol. Gen. d'It., 26a C.G.I., Paris 80, G13, p. 65-70.
- Castellarin A., 1981. *Carta tettonica delle Alpi Meridionali*. Alla scala 1:200000., C.N.R. Progetto Fin. Geod., Sottoprogetto 5 "Modello Strutturale". Pubbl. n. 441, p. 1-217.
- Castellarin A., Colacicchi R., Praturlon A., 1978. *Fasi distensive, trascorrenze e sovrascorrimenti lungo la "Linea Ancona-Anzio", dal Lias Medio al Pliocene*. Prog. Fin. Geod., Pubbl. N. 89, Geol. Rom. 17, p. 1-26, Pos. B3.
- Castellarin A., Frascari F., Val G.B., 1979. *Problemi di interpretazione geologica profonda del sudalpino orientale*. Rend. Soc. Geol. It., v. 2, p. 55-60.
- Castellarin A., Val G.B., 1981. *Importance of Mercynian Tectonics within the framework of the Southern Alps*. Journal of Structural Geology, vol. 3, n. 4, pp. 477-486.
- Catalano R., D'Argenio B., 1978. *An essay of planispastic restoration across Western Sicily*. Estratto da Geologica Romana, v. XVII, p. 145-159.
- Channel J.E.T., D'Argenio B., Horvath F., 1979. *Adria, the African Promontory, in Mesozoic Mediterranean Palaeogeography*. Earth Science Reviews, p. 213-292.
- Cherchi A., 1973. *Appunti biostratigrafici sul Pliocene in Sardegna*. Boll. Soc. Geol. It., 92 (1973), pp. 891-902.
- Chumakov I.S., 1968. *Pliocene ingression in the Nile Valley according to new data*. In: Desert and River in Nubia, Geomorphology and Prehistoric Environments at the Aswan Reservoir. Butzer K.A., and Hansen C.L., eds., Madison: The University of Wisconsin Press, p. 521-522.
- Cita M.B., Colalongo M.L., D'Onofrio S., Iaccarino S., Salvatorini G., 1978. *Biostratigraphy of Miocene Deep Sea Sediments (Sites 372 and 375), with special Reference to the Messinian/Pre-Messinian Interval*. Initial Reports of the Deep Sea Drilling Project, p. 671-685.
- Cita M.B., Ryan W.B., 1973. *The Pliocene record in deep sea Mediterranean sediments. Time-scale and general synthesis*. In: Ryan W.B.F., HSU K.J., et al., Initial Reports of the Deep Sea Drilling Project. Vol. 13: Washington (U.S. Government Printing Office), p. 1405-1415.
- Cita M.B., Wright R.C., Ryan W.B.F., Longinelli A., 1978. *Messinian paleoenvironments*. Initial Reports of the Deep Sea Drilling Project, v. 42, Part. 1, p. 1003-1032.
- Cloeting, Sierd, Nolet G., Wortel R., 1980. *Crustal structure of the eastern Mediterranean inferred from Rayleigh wave dispersion*. E.P.S.L., 51, p. 336-342.
- Closs H., Hinz K., 1974. *Seismische und bathymetrische Ergebnisse von Mediterranean Rucken und Hellenischen Graben*. Aeg. Symposium, Hannover 1974.
- Colacicchi R., Piali P., Praturlon A., 1975B. *Megabreccias as a product of tectonic activity along a Carbonate platform margin*. IX. Int. Sediment Congr. Theme 4, no. 1: 61-66.
- Colantoni P., Borsetti A.M., 1973. *Some notes on geology and stratigraphy of the Strait of Sicily*. Bull. Geol. Soc. Greece, v. 10, p. 31-32.
- Colantoni P., Zarudski E.F.K., 1972. *Some principal seafloor features in the Strait of Sicily*. 23e Congrès C.I.E.S.M., Athènes, 3-11 Novembre 1972. Comité de géologie et de géophysique marines, 3 pp.
- Colom G., Escandell B., 1962. *L'évolution du Géosynclinal Baléare*. Soc. Géol. France, Livre P. Falot, v. 1, p. 125-136.
- Cominakis P.E., Papazachos D.C., 1972. *Seismicity of the eastern Mediterranean and some tectonic features of the Mediterranean Ridge*. Bull. Geol. Soc. Amer., v. 83, p. 1093-1102.
- Conant L.C., Goudarzi G.H., 1967. *Stratigraphic and tectonic framework of Libya*. Bull. Amer. Ass. Petrol. Geol., v. 51 (5), p. 719-730.
- Cortésogno L., Galbiati B., Principi G., 1979. *Le breccie serpentinitiche giurassiche della Liguria Orientale*. Archiv. de Sc., 33: 185-200. Special issue on Tectonic inclusions and associated rocks in serpentinites. Geneva, September, 1979.
- Coulon C., Bobier C., Demant A., 1973. *Contribution du paléomagnétisme à l'étude des séries volcaniques cénozoïques et quaternaires du Logudoro et du Bosano (Sardaigne Nord-occidentale)*. Le problème de la dérivée de la Sardaigne, Monaco: CIESM, Rapport Scientifique No. 6, p. 59-61.
- Coulon C., Demant A., Bobier C., 1974b. *Contribution du paléomagnétisme à l'étude des séries volcaniques Cénozoïques et quaternaires de Sardaigne Nord-occidentale*. Tectonophysics, v. 22, p. 59-82.
- Cruz, J.Y. and Rapp, R.H., 1982. *Sea Surface in the Mediterranean Area from SEASAT Altimeter Data*. Boll. Geof. Teor. Appl. Vol. 24, No. 95.
- Dalipi H., Kondo A., Pejo I., Ikonomi J., Mecaj B., 1971. *Stratigraphy of the Deposits of the Mesozoic in Southern and Western Albania (outer Albanids)*. Nafta, Zagreb, v. 4-5 (XXII), p. 227-253.
- D'Argenio B., Radoicic R., Sgossio I., 1971. *A paleogeographic section through the Italo-Dinaric external zones during Jurassic and Cretaceous times*. Symposium on exploration for oil and gas deposits in the Adriatic Sea and Outer Dinarides zones. Zara, 1971.
- Dewey J.F., Bird J.M., 1970. *Mountain belts and the new global tectonics*. J. Geophys. Res., v. 75, p. 2625-2647.
- Dewey J.F., Pitman W.C., Ryan W.B.F., Bonnin J., 1973. *Plate tectonics and the evolution of the Alpine system*. Geol. Soc. Amer. Bull., v. 84, p. 3137-3180.
- Desmons J., 1977a. *Ophiolites and melange in the Maskutan-Fanuj area and south-east of Iranshar (eastern Baluchistan, Iran)*. Geol. Surv. Iran, unpubl. Rept., 25 pp.
- Dietrich V., Emmertmann R., Puchelt H. Keller J., 1978. *Oceanic basalts from the Tyrrhenian Basin*. DSDP Leg. 42A, Hole 373A., Initial Reports of the Deep Sea Drilling Project, v. 42, p. 515-530.
- Di Paola G., 1973. *The Island of Linosa (Sicily Channel)*. Bull. Volcanol. v. 37, n. 2.
- Druckman Y., Gvirtzman G., 1975. *Distribution and environments of deposition of Upper Triassic on the northern margins of the Arabian shield and around the Mediterranean sea*. IXème Congrès Internat. Séd. Thème 5 p. 183-189.
- Dubertret L., 1970. *Revue of the structural geology of the Red Sea and the surrounding areas*. Phil. Trans. Roy. Soc. London Ser. A, v. 267, p. 9-20.
- Dubertret L., 1976. *La péninsule Arabique: explanatory text to sheet 16. Tectonic map of Europe 1:2.500.000, Chapter 6, Section 4 I.G.C. Sub. Commission of Tectonic Maps*.
- El Adl R., El Naggag G., 1975. *Geothermal gradient map of the Gulf of Suez region*. Egypt, Unpublished

- map. General Petroleum Company, Cairo.
- Elter P., 1973. *Lineamenti tettonici ed evolutivi dell'Appennino settentrionale*. In: *Moderne vedute sulla Geologia dell'Appennino*. Roma: Acad. Naz. Lincei, p. 97-118.
- Emery K.O., Heezen B.C., Allan T.D., 1966. *Bathymetry of the eastern Mediterranean Sea*. Deep-Sea Res., v. 13, p. 173-192.
- Erickson A., J., 1970. *The measurements and interpretation of heat flow in the Mediterranean and Black Seas*. Ph. D. Thesis, Cambridge, Mass.: Mass. Inst. Technology, 272 pp.
- Erickson A.J., Simmons G., Ryan W.B.F., 1976. *Review of Heat Flow data from the Mediterranean and Aegean Seas*. In: *The structural History of the Mediterranean Basins*. (B. Bijou-Duval and L. Montadert, edn.). Edition Technip.
- Erickson A.J., Von Herzen R.P., 1978. *Down-hole temperature measurements*. Deep Sea Drilling Project, Leg. 42A, v. XLII, Part. I, p. 857-871.
- Fabbri A., Curzi P., 1978. *A seismic reflection study of the Messinian event in the Tyrrhenian sea*. Paper presented to Internat. Meeting on Geodynamic and Biodynamic effects of Messinian Salinity Crisis in the Mediterranean, Accad. Naz. Lincei, Rome.
- Fabbri A., Selli R., 1972. *The structure and stratigraphy of the Tyrrhenian Sea*. In: *The Mediterranean Sea — A Natural Sedimentation Laboratory*. Stanley, D.J., ed., stroudsburg: Dowden, Hutchinson, Ross, p. 75-81.
- Fabrics F., Schmidt-Thome P., 1972. *Contribution to recent sedimentation on the shelves of the Southern Adriatic, Ionian and Syrtis seas*. In: *The Mediterranean Sea — A natural Sedimentation Laboratory*. Stanley D.J., Stroudsburg: Dowden, Hutchinson, Ross, p. 333-343.
- Fahlquist D.A., Hersey J.B., 1969. *Seismic refraction measurements in the western Mediterranean Sea*. Bull. Inst. Océanogr. Monaco, v. 67 (1386) 52 pp.
- Felix R., 1973. *Oligo-Miocene stratigraphy of Malta and Gozo*. 1-104, Meded. Landbown. Megeningen.
- Finetti I., 1976. *Mediterranean Ridge. A young submerged chain associated with the Hellenic arc*. Bull. Geol. Teor. Appl. v. 28, N. 69, p. 31-62.
- Finetti I., 1981. *Mediterranean Sea: one of the world's most promising deep water oil basins*. Deep offshore technology conference proceedings, v. 1, p. 12-23.
- Finetti I., 1981. *Geophysical study, on the evolution of the Ionian Sea*. Sedimentary basins of Mediterranean margins Ist. di Geol. Urbino. C.N.R. Italian Project of Oceanography, p. 465-484.
- Finetti I., Belleo S., De Visintini G., 1966. *Preliminary investigation on the Earth's crust in the South Adriatic Sea*. Boll. Geol. Teor. Appl. VIII (29): 2-39.
- Finetti I., Morelli C., 1972. *Wide scale digital seismic exploration of the Mediterranean Sea*. Geofis. Teor. Appl., v. 14, p. 291-342.
- Finetti I., Morelli C., 1973. *Geophysical exploration of the Mediterranean Sea*. Boll. Geofis. Appl., v. 15, p. 263-341.
- Finetti I., Morelli C., Zarudski E., 1970. *Reflection seismic study of the Tyrrhenian Sea*. Boll. Geofis. Teor. Appl., v. 12, p. 311-346.
- Foose R.M., Brigham R.M., 1976. *Interpreting the genetic character of structural lineaments in the Mediterranean basin*. Proceedings of the Second International Conference on Basement Tectonics, Newark, Delaware, b. 211-222.
- Galdeano A., Courtillot V., Le Borgne E., Le Mouel J.L., Rossignol J.C., 1974. *Aeromagnetic survey of the Mediterranean: description and tectonic implications*. Earth Planet. Sci. Lett., v. 23 (3), p. 323-336.
- Garrison R.E., Schreiber B.C., Bernoulli D., Fabricius F.H., Kidd R.B., Mélières F., 1978. *Sedimentary Petrology and Structures of Messinian Evaporitic Sediments in the Mediterranean Sea*. Leg. 42A, Deep Sea Drilling Project, Initial Reports of the Deep Sea Drilling Project., v. 17, p. 571-611.
- Gaskell J.F., Swallow J.C., 1953. *Seismic refraction experiments in the Mediterranean and Indian Ocean*. Nature, v. 172, p. 535.
- Gasparini C., Iannaccone G., Scandone P., Scarpa R., 1982. *Seismotectonic of the Calabrian Arc*. Tectonophysics, 83, Tec. 383.
- Gass I.G., 1968. *Is the Troodos massif of Cyprus a fragment of Mesozoic ocean floor?* Nature, v. 220, p. 39-42.
- Gass I.G., Smewing J.D., 1973. *Intrusion, extrusion and metamorphism at constructive margins: evidence from the Troodos Massif, Cyprus*. Nature, v. 242, p. 26-29.
- Gergawi A., Khashab H., 1968. *Seismicity of the U.A.R.* Bull. Helwan Observ., v. 76, 27.
- Giorgetti F., Iaccarino E., 1971. *Seismicity of the Italian region*. Boll. Geofis. Teor. Appl., v. 13 (50), p. 143-154.
- Giunta G., Liguori V., 1972. *Geologia dell'estremità Nord-Occidentale della Sicilia*. Rivista Mineraria Siciliana. Anno XXIII, n. 136-138, p. 165-226.
- Glangeaud L., 1962. *Paléogéographie dynamique de la Méditerranée et de ses bordures. Le rôle des phases ponto-plio-quaternaires*. In: *Océanographie, Géologie et Géophysique de la Méditerranée Occidentale*. le. Bourcart J., ed., Coll. Nat. Cent. Nat. Rech. Sci., Villefranche-sur-Mer, p. 125-165.
- Geodick T.R., Sagabiel S., 1976. *Sediment movement in Lebanese submarine canyons*. In: *Symposium on the Eastern Mediterranean*. Hulings N., ed., Acta Adriatica, Split, v. 18, p. 117-128.
- Goncharov V.P., Mikhaylov O.V., 1964. *New data concerning the bottom relief of the Mediterranean*. Deep Sea Res., v. 11, p. 625.
- Gonnard R., Letouzey J., Bijou-Duval B., Mointadert L., 1975. *Apports de la sismique réflexion aux problèmes du volcanisme en Méditerranée et à l'interprétation des données magnétique*. Science de la Terre. Montpellier, Abstract.
- Got H., Stanley D.J., Sorel D., 1977. *Northwestern Hellenic Arc: concurrent sedimentation and deformation in a compressive setting*. Mar. Geol., v. 24, p. 21-36.
- Grandjacquet C., 1963. *Schéma structural de l'Apennin Campano-lucanien (Italie)*. Rev. Géogr. Phys. Géol. Dyn., Paris, v. 5, p. 185-202.
- Grandjacquet C., Haccard D., Lorenz C., 1972. *Essai de tableau synthétique des principaux événements affectant les domaines alpin et apennin à partir du Trias*. C.R. Somm. S.G.F., p. 158-163.
- Gregor C.B., Nairn A.E.M., 1971. *Paleomagnetic results from Lebanon (abstract)*. Trans. Amer. Geophys. Union, v. 52 (4), p. 188.
- Hamilton N., Hailwood E.A., Kidd A.B., 1978. *Preliminary paleomagnetic chronology of Cenozoic sediments from DSDP site 372, 374, and 376 of the Mediterranean Sea*. Initial Reports of the Deep Sea Drilling Project, v. 42, Part. 1, p. 873-879.
- Hammuda O.S., 1969. *Jurassic and Lower Cretaceous Rocks of Central Jebel Nefusa, Northwest Libya*. Petrol. Explor. Soc. Libya, 74.
- Hammuda O.S., 1971. *Nature and significance of the Lower Cretaceous unconformity in Jebel Nefusa, Northwest Libya*. In: *Symp. Geol. Libya* (ed. C. Gray), Fac. Sci., Univ. Libya, Tripoli, 87-97.
- Hammuda O.S., Missallati A.A., 1980. *A study of the Libyan-Tunisian continental shelf*. Technical Report submitted to the International Court of Justice, The Hague.
- Harrison J.C., 1955. *An interpretation of gravity anomalies in the Eastern Mediterranean*. Phil. Trans. Roy. Soc. London, v. A. 248, p. 283-325.
- Hassan A.H., 1962. *Distribution of Triassic and Jurassic formations in east Mediterranean*. Proc. 4th Arab. Petrol. Congr. (Beirut), Paper 25 (B3).
- Heimann K.O., 1975. *Comparison of late Miocene Strata in Sicily and on the Ionian Islands*. Rapp. comm. in. Mer Médit., 23, 4a, 147.
- Hersey J.B., 1965. *Sedimentary basins of the Mediterranean Sea*. In *Submarine Geology and Geophysics*. Whittard W.F., Bradshaw R., eds., London: Butterworths, Colston, Paper 17, p. 75-91.
- Hieke W., 1976. *Problems of eastern Mediterranean late Quaternary stratigraphy — a critical evaluation of literature*. Meteor. Forschungs-Ergebnisse, v. 24, p. 68-88.
- Hinz K., 1974. *Results of seismic refraction and seismic reflection measurements in the Ionian Sea*. Geol. Jb., v. E-2, p. 33-65.
- Horvath F., Channell T.E.T., 1977. *Further evidence relevant to the African/Adriatic Promontory as a Paleogeographic Premise for Alpine Orogeny*. In: (B. Bijou-Duval and L. Montadert, eds.). *International Symposium on the Structural History of the Mediterranean Basin, Split, (Yugoslavia)*, Ed. Technip., Paris, p. 133-142.
- House M.R., Dunham K.C., Wigglesworth J.C., 1962. *Geology and Structure of the Maltese Islands*. In: Bowen-Jones H., et al., Malta, 25-33, Newcastle upon Tyne.
- Hsu J.K., 1978. *Genesis of the Tethys and the Mediterranean*. Initial Reports of the Deep Sea Drilling Project, v. 42, Part. 1, p. 943-948.
- Hsu K.J., Ryan W.B.F., 1973. *Deep-Sea drilling in the Hellenic Trench*. Geol. Soc. Greece Bull., v. 10 (1), p. 81-80.
- Hurley R.J., 1971. *Seismic reflection studies in the Adriatic Sea*. Rev. Géogr. Phys. Géol. Dyn., v. 13 (5), p. 429-438.
- Hynes A.J., Nisbet E.G., Smith A.G., Welland M.J.P., Rex D.C., 1972. *Spreading and emplacement ages of some ophiolites in the Othris region, eastern central Greece*. Zeitschr. Deutsch. Geol. Ges., v. 123, p. 445-468.
- Ilhan E., 1945. *Le volcanisme néogène et quaternaire Anatolie*. Turk. Congr. Dergisi (Ankara), v. 3, p. 1-8.
- Ilhan E., 1967. *Toros-Zagros folding and its relation to the Middle East oil field*. Bull. Amer. Assoc. Petrol. Geol., v. 51 (5), p. 651-667.
- Ilhan E., 1971. *The structural features of Turkey*. In: *Geology and History of Turkey*, Campbell A.S., ed., Tripoli: Petrol Explor. Soc. Libya, p. 159-170.
- Ippolito F., D'Argenio B., Pescatore T., Scandone P., 1975. *Structural stratigraphic units and tectonic framework of southern Appennines*. Reprinted from *Geology of Italy*. Coy Squyres, ed., The Earth Sciences Soc. of the Libyan Arab Republic, Tripoli, p. 317-328.
- Jong K.A. de, Manzoni M., Stavenga T., van der Voo R., van Dijk F., Zijdeveld D.A., 1972. *Early Miocene*

- age of rotation of Sardinia: paleomagnetic evidence. 23e Congrès-Assemblée, C.I.E.S.M., Athènes, 3-11 Novembre 1972, 3 p.
- Jong K.A. de, van der Voo R., 1970. Rotation of Sardinia: paleomagnetic evidence from Permian rocks. *Nature*, v. 226, p. 933-934.
- Jongsma D., 1974. Heat flow in the Aegean Sea. *Geophys. J.R. Astr. Soc.*, v. 37, p. 337-346.
- Jongsma D., Wissman G., et al., 1975. The southern Aegean Sea: an extensional marginal basin without spreading? *Bundes. Geowissenschaften Rohstoffe*, 41 pp.
- Kaya O., 1975. Northwest Anatolian ophiolite zone (Nonmetamorphic and Mesozoic in age). (Abstract), Orsay, France: Vème Colloque sur la Géologie des Régions Egéennes.
- Kazmin V.G., Kulakov V.V., 1965. Formation of ophiolites in the Northwest of Syria. *Izvest. Vys. Ucheb. Zaved. Geol. y Razv.*, v. 2, p. 3-14 (in Russian).
- Kazmin V.G., Kulakov V.V., 1968. The Geological Map of Syria, Scale 1:500,000; Explanatory Notes (for Part of the Lattaqiah Region). Damascus: Syrian Arab Republic, Ministry of Petroleum, Electricity and Execution of Industrial Projects, 124 p.
- Keller J., Ryan W.B.F., Ninkovich D., Altherr R., 1974. The deep sea record of Quaternary volcanism in the Mediterranean. XXIVe Congrès-Assemblées plénières de Monaco, 6-14 Décembre 1974, Comité Géol. Géophys. Mar., 4 pp.
- Kenyon N.H., Belderson R.H., 1977. Young compressional structures of the Calabrian, Hellenic and Cyprus outer ridges. In: Structural History of the Mediterranean Basins. Technip, ed., Paris, p. 233-240.
- Khan M.A., Summers C., Bamford S.A.D., Chroston P.N., Poster C.K., Vine F.J., 1972. A reversed seismic refraction line on the Troodos Massif, Cyprus. *Nature Phys. Sci.*, v. 238, no. 87, p. 134-136.
- Klitzsch E., 1968. Outline of the Geology of Libya. In: Geology and Archaeology of Northern Cyrenaica, Libya, 10th Annual Field Conference, Petr. Expl. Soc. of Libya, p. 71-78.
- Klootwijk C.T., van den Berg J., 1975. The rotation of Italy: Preliminary palaeomagnetic data from the Umbrian sequence, Northern Apennines. *Earth Planet. Sci. Lett.*, v. 25, p. 263-273.
- Kondo A., 1970. De la présence de structures ioniennes à partir du Jurassique inférieur et de leur développement sous-marin. *Bull. U.S.H.T.*, Tirana, n. 4.
- Lanzafame G., Spadea P., Tortorici L., 1976. Serie ofiolitifere nell'area fra Tarsia e Spezzano Albanese (Calabria): Stratigrafia, Petrografia, Rapporti Strutturali. C.N.R., Ofioliti, Anno I N. 3, p. 464-467.
- Laubscher H.P., 1970. Bewegung und Wärme in der alpinen Orogenese. *Schweiz. Min. Petrogr. Mitt.*, v. 50, p. 503-534.
- Laubscher H.P., 1970. Das Alpen-Dinariden-Problem und die Palinopastik der Südlichen Tethys. *Geol. Rundsch.*, v. 60, p. 813-833.
- Laubscher H.P., 1974. Evoluzione e struttura delle Alpi. *Le Scienze*, v. 72 p. 48-59.
- Laubscher H.P., 1975. Plate boundaries and microplates in Alpine history. *Amer. J. Sci.*, v. 275, p. 865-876.
- Laubscher H.P., Bernouilli D., 1977. Mediterranean and Tethys, the Ocean Basins and Margins. The Eastern Mediterranean, v. 4A, p. 1-22.
- Le Borgne E., Bayer R., Le Mouel J.L., 1972. La cartographie magnétique de la partie sud du bassin algéro-provençal. *Compt. Rend.*, v. 274, p. 1291-1294.
- Le Houel J., Le Borgne E., 1970. Les anomalies magnétiques du Sud-est de la France et de la Méditerranée Occidentale. *C.R. Acad. Sci. Paris, Sér. D.*, v. 271, p. 1348-1350.
- Le Borgne E., Le Mouel J.L., Le Pichon W., 1971. Aeromagnetic survey of southwestern Europe. *Earth Planet. Sci. Lett.*, v. 12, p. 287-299.
- Le Pichon X., Angelier J., 1979. The Hellenic arc and trench system: A key to the neotectonic evolution of the eastern Mediterranean area. *Tectonophysics*, v. 60, p. 1-42.
- Le Pichon X., Angelier J., Auboin J., Lyberis N., Monti S., Renard V., Got H., Hsu K.J., Mart Y., Mascle J., Matthews D., Mitropoulos D., Tsoflias P., Chronis C., 1979. From subduction to transform motion. A seabeam survey of the Hellenic trench system. *Earth and Planetary Science Letters*, v. 44, p. 441-450.
- Letouzey J., Tremolieres P. and B., Bijou-Duval B., 1977. An approach to the Structure of the Mediterranean Area: A Satellite Photogeological Study. In: B. Bijou-Duval and L. Montadert (eds.), Structural History of the Mediterranean Basins, Edition Technip., p. 215-219.
- Lort J.M., 1972. The Crustal Structure of the Eastern Mediterranean. Ph. D. Thesis, Univ. Cambridge, Cambridge, 117 pp.
- Lort J.M., 1973. Summary of seismic studies in the eastern Mediterranean. *Bull. Geol. Soc. Greece*, v. 10, p. 99-108.
- Lort J.M., 1977. Geophysics of the Mediterranean Sea Basins. In: Nairn A.E.M., Kanes W.H., Stehli F.G. — The Ocean Basins and Margins, 4A. The Eastern Mediterranean, 151-213, New York.
- Lort J.M., Gray F., 1974. Cyprus: seismic studies at sea. *Nature*, v. 248, p. 745-747.
- Lort J.M., Limond W.Q., Gray F., 1974. Preliminary seismic studies in the eastern Mediterranean. *Earth. Planet. Sci. Lett.*, v. 21, p. 355-366.

- Lowrie W., Alvarez W., 1975. Paleomagnetic evidence for rotation of the Italian Peninsula. *J. Geophys. Res.*, v. 80, p. 1579-1592.
- Makris J., 1975. Crustal structure of the Aegean Sea and the Hellenides obtained from geophysical survey. *J. Geophys.*, v. 41, p. 441-443.
- Maldonado A., Stanley D.J., 1976a. Late Quaternary sedimentation and stratigraphy in the strait of Sicily. *Smithson. Contrib. Earth Sci.*, v. 16, 73 pp.
- Maldonado A., Stanley D.J., 1976b. The Nile Cone: submarine fan development by cyclic sedimentation. *Mar. Geol.*, v. 20, p. 27-40.
- Malovitskij J.P., Chumakov, I.S., Shimkus, K.M., Esina, P.A., Moskalenko, V.N., editors, 1982. The Earth's Crust and the History of Evolution of the Mediterranean Sea. Acad. Sciences U.S.S.R., Nauka ed., 1982.
- Malovitskij J.P., Emelyanov E.M., Kaazakov O.V., Moskalenko V.N., Osipov G.V., Shimkus K.M., Chumakov I.S., 1975. Geological structure of the Mediterranean Sea floor (based on geological-geophysical data). *Mar. Geol.*, v. 18, p. 231-261.
- Marini M., 1980. La linea Sestri-Voltaggio nel quadro dell'evoluzione tettonogenetica del limite Alpi-Appennino: proposta di interpretazione. *Mem. Soc. Geol. Ital.*, 19 (1978): 445-452.
- Martin L., 1967. Geology and History of Libya. *Petrol. Expl. Soc.*, volume Amsterdam: Breumelhof N.V.
- Martin L., Bobier G., 1977. La distension cenozoïque du détroit siculo-tunisien, evolution des contraintes dans un rift naissant en milieu décontinental. 5° Reun. Ann. Sc. Terre, Rennes, 19-22 Avril 1977, p. 327.
- Martinis B., Pieri M., 1963. Alcune notizie sulle formazioni evaporitiche del Triassico superiore nell'Italia centrale e meridionale. *Mem. Soc. Geol. Ital.*, v. 4, 30 p.
- Marussi, A., 1980. Geoidal features and crustal structure in Central Asia. *Proc. Qinghai-Xizang (Tibet) Plateau Symposium*, Beijing, 1980.
- Mascle G., 1974. Les grands traits de l'évolution géologique des monts Sicani (Sicile). *Bull. Soc. Géol. Fr.* (7), 16, 161-170.
- Matthews D.H., 1974. Bathymetry and magnetic anomalies of the Eastern Mediterranean. *CIESM Congr.*, December, 1974, Monaco, Abstract.
- Mercier J.L., Carey E., Philip H., Sorel D., 1976. La néotectonique plio-quaternaire de l'arc égéen externe et de la mer Egée et ses relations avec la sismicité. *Bull. Soc. Géol. France*, v. 13, p. 367-384.
- Mikbel S.R., 1979. Structural and configuration map of the basement of east and central Libya. *N.J. Geol. Paläont. Ab.*, 158, 2, p. 203-220.
- Mikbel S.R., 1977. Basement configuration and structure of West Libya. *The Libyan Journal of Science*, v. 7A, p. 19-34.
- Missallati A.A., 1980. Hydrocarbon productive trends in and adjacent to the Pelagian basin. Technical Report submitted to the International Court of Justice, The Hague.
- Missallati A.A., Hammuda O.S., et al., 1979. A study of the Libyan-Tunisian Continental Shelf. Unpubl. Technical Report, No. 1.
- Miyashiro A., 1973. The Troodos ophiolite complex was probably formed in an island arc. *Earth Planet. Sci. Lett.*, v. 19, p. 218-224.
- Montadert L., Letouzey J., Mauffret A., 1978. Messinian Event. Seismic Evidence. Initial Reports of the Deep Sea Drilling Project, v. 42, Part 1, p. 1037-1048.
- Moore E.M., 1969. Petrology and structure of the Vourinos ophiolite complex, northern Greece. *Geol. Soc. Amer. Spec. Pap.*, no. 118, p. 1-73.
- Moore E.M., Vine F.J., 1971. The Troodos Massif, Cyprus and other ophiolites as ocean crusts: evaluation and implications. *Roy. Soc. London Phil. Trans. (A)*, v. 268, p. 443-446.
- Morelli C., 1970. Physiography, gravity and magnetism of the Tyrrhenian Sea. *Boll. Geofis. Teor. Appl.*, v. 12, p. 274-308.
- Morelli C., 1972. Bathymetry, gravity and magnetism in the Strait of Sicily. In: *Oceanography in the Strait of Sicily*. Allan T.D., Akal T., Malcard R., eds., Saclant ASW Res. Cent., La Spezia, Conf. Proc. 7, p. 193-207.
- Morelli C., Pisani M., Gantar C., 1975. Geophysical studies in the Aegean Sea and in the eastern Mediterranean. *Boll. Geof. Teor. Appl.*, v. 18, p. 127-167.
- Morgan P., Evans G., 1975. Recent tectonics of the Cilicia Plateau and the north-eastern Mediterranean. *EOS*, v. 56 (12), p. 1058.
- Moskalenko V.N., 1965. Study of sedimentary series of the Mediterranean Sea by seismic methods. In: Basic Features of the Geological Structure of the Hydrologic Regime and Biology of the Mediterranean Sea. Translation and Interpretation Division, Institute of Modern Languages, p. 60-72.
- Moskalenko V.N., 1966. New data and structure of sedimentary strata and basement in the Levant Sea. *Oceanology*, v. 6, p. 828-836.
- Moskalenko V.N., 1967. Study of the sedimentary series of the Mediterranean Sea by Seismic methods. *Proc.*

- Acad. Sci. USSR Oc. Comm. (Transl. 1969).
- Moskalenko V.N., Yelnikov I.N., 1966. *Seismic data about the possible continuation of the African Platform in the Crete-African region of the Mediterranean Sea*. Int. Oceanogr. Congr., v. 2, p. 259-260.
- Mulder C.J., 1973. *Tectonic framework and distribution of Miocene evaporites in the Mediterranean*. In: Messinian Events in the Mediterranean. Drooger C.W., ed., Amsterdam: North Holland Publ. Co., p. 44-59.
- Muller J., Hieke W., Fabricius F., 1978. *Turbidites at site 374: their composition, provenance and paleobathymetric significance*. Initial Reports of the Deep Sea Drilling Project, v. 42, p. 397-400.
- Mur A., Ben Avraham Z., 1978. *The eastern Mediterranean and the Levant: Tectonics of continental collision*. Tectonophysics, v. 46, p. 297-311.
- Nardi R., Dallan L., 1979. *Osservazioni sulle fasi tettoniche tra l'Oligocene e il Miocene Inferiore nell'Appennino Settentrionale*. C.N.R., Prog. Fin. Geod., Sottoprogr., Modello Strutturale, Riunione "Profili Geologici attraverso la Penisola Italiana". Pisa, aprile 1978, Rapp. Int. N. 165, pp. 14-18.
- Neev D., 1975. *Tectonic evolution of the Middle East and the Levantine Basin (easternmost Mediterranean)*. Geology, v. 3, p. 683-686.
- Nesteroff W.D., 1972. *Mineralogy, petrography, distribution, and origin of the Messinian Mediterranean Evaporites*. In: Ryan W.B.F., Hsu K.J., et al., Initial Reports of the Deep Sea Drilling Project, Washington D.C.: U.S. Government Printing Office, v. 13, p. 673-694.
- Ogniben L., Martinis B., Rossi P.M., et al., 1973. *Structural model of Italy, North and South; lithostratigraphic-tectonic map, bathymetric map*. C.N.R., Map.
- Ozgul N., Arpat E., 1973. *Structural units of the Taurus orogenic belt and their continuation in neighboring regions*. Geol. Soc. Greece Bull., v. 10, p. 156-164.
- Papa A., 1970. *Conceptions nouvelles sur la structure des Albanides (présentation de la carte tectonique de l'Albanie au 500,000^e)*. Bull. Soc. Géol. France, v. 12, p. 1096-1109.
- Papazachos B.C., 1973. *Distribution of seismic foci in the Mediterranean and surrounding area and its tectonic implication*. Geophys. J.R. Astr. Soc., v. 33, p. 421-430.
- Papazachos B.C., Comninakis P.E., 1978. *Deep Structure and tectonics of the eastern Mediterranean*. Tectonophysics, 46, p. 285-296.
- Patacca E., Scandone P., Giunta G., Liguori V., 1979. *Mesozoic Paleotectonic evolution of the Ragusa zone (Southeastern Sicily)*. Estratto da: Geologia Romana, v. 18, p. 331-369.
- Pedley H.M., Ghouse M.R., Waugh B., 1978. *The Geology of the Pelagian Block: the Maltese Islands, the Ocean Basins and Margins*. In: The Western Mediterranean, v. 4B, p. 417-435.
- Pieri M., Groppi G., 1981. *Subsurface geological structure of the Po Plain, Italy*. C.N.R. Prog. Fin. Geodinamica, AGIP, Pubbl., N. 414.
- Pinar N., 1960. *La carte seismotectonique de la Turquie*. U.G.G.I., Monogr., no. 6.
- Pinar-Erdem N., 1974. *The Turkish green rocks*. Bull. Miner. Res. Explor. Inst. Turk., v. 8, p. 119-131.
- Pinar N., Ilhan E., 1977. *Outlines of the stratigraphy and tectonics of Turkey, with notes on the Geology of Cyprus, the Ocean basins and margins*. The Eastern Mediterranean, v. 4A, p. 277-313.
- Pitman W.C., Talwani M., 1972. *Sea-floor spreading in the North Atlantic*. Geol. Soc. Amer. Bull., v. 83, p. 619-646.
- Poizat C., 1970. *Hydrodynamisme et sédimentation dans le Golfe de Gabès (Tunisie)*. Téthys, v. 2, p. 267-296.
- Ponikarov V.P., Kazmin V.G., Mikhailov I.A., Razvaliyev A.V., Krasheninnikov V.A., Kozlov V.V., Souliidi-Kandratyev E.D., Mikhailov K.Y., Kulakov V.V., Faradzhev V.A., Mirzayev K.M., 1967. *The Geology of Syria: Explanatory Notes on the Geological Map of Syria, Scale 1:500,000, Part I, Stratigraphy, Igneous Rocks and Tectonics*. Damascus: Syrian Arab Republic, Ministry of Industry, 229 p.
- Rabinowitz P.D., Ryan W.B.F., 1970. *Gravity anomalies and crustal shortening in the eastern Mediterranean*. Tectonophysics, v. 10, p. 585-608.
- Rampnoux J.P., 1973. *Essai de reconstitution géotectonique des Dinarides internes yougoslave (Serbie) au Jurassique et au Crétacé*. Réunion Annu. Sci. de la Terre, Paris, p. 353.
- Richert J.P., 1971. *Mise en évidence de quatre phases tectoniques successives en Tunisie*. Notes Serv. Geol. Tunisie, no. 34, p. 115-125.
- Ricou L.E., 1971. *Le croissant ophiolitique péri-arabe, une ceinture de nappes mises en place au Crétacé supérieur*. Rev. Géogr. Phys. Géol. Dyn., v. 13, p. 327-349.
- Ricou L.E., Argyriadis I., Lefevre R., 1974. *Proposition d'une origine interne pour les nappes d'Antalya et le massif d'Alanya (Taurides occidentales, Turquie)*. Soc. Géol. France Bull., v(7) 16, p. 107-111.
- Robertson A.H.F., Hudson J.D., 1974. *Pelagic sediments in the Cretaceous and Tertiary history of the Troodos Massif, Cyprus*. Int. Ass. Sediment., Spec. Publ. no. 1, p. 403-436.
- Ryan W.B.F., 1972. *Stratigraphy of Late Quaternary sediments in the eastern Mediterranean*. In: The Mediterranean Sea — A Natural Sedimentation Laboratory. Stanley D.J., ed., Stroudsburg: Dowden,

- Hutchinson, and Ross, p. 149-169.
- Ryan W.B.F., 1973. *Geodynamic implications of the Messinian crisis of salinity*. In: Messinian Events in the Mediterranean. Drooger C.W., ed., Amsterdam: North Holland Publ. Co., p. 26-38.
- Ryan W.B.F., 1978. *Messinian badlands on the southeastern margin of the Mediterranean Sea*. Mar. Geol., 27: 349-363.
- Ryan W.B.F., Cita M.B., 1978. *The nature and distribution of Messinian erosional surfaces — indicators of a several-kilometer-deep Mediterranean in the Miocene*. Marine Geology, v. 27, p. 193-230.
- Ryan W.B.F., Ewing M. and Ewing T.I., 1966. *Diapirism in the sedimentary basins of the Mediterranean Sea*. Trans. Amer. Geophys. Union, v. 47, p. 120.
- Ryan W.B.F., Stanley D.J., et al., 1970. *The tectonics and geology of the Mediterranean Sea*. In: The Sea. Maxwell A.E., ed., New York: Wiley-Interscience, v. 4 (part. 2), p. 387-492.
- Said R., 1962. *The Geology of Egypt*. Amsterdam: Elsevier Publ. Co., 377 pp.
- Sander N.J., 1970. *Structural evolution of the Mediterranean Region during the Mesozoic Era*. In: Geology and History of Lybia, Alvarez, Gohrbandt K., eds., Tripoli: Petr. Exploration Soc. Libya, p. 43-132.
- Scandone P., 1975. *The seaways and the Jurassic Tethys Ocean in the central Mediterranean area*. Nature, v. 256, no. 5513, p. 117-119.
- Scandone P., 1978. *Origin of the Tyrrhenian Sea and Calabrian Arc*. Boll. Soc. Geol. It., p. 27-34.
- Scandone P., Giunta G., Liguori V., 1974. *The connection between the Apulia and Sahara continental margins in the southern Apennines and in Sicily*. Cong. Ass. CIESM, Monaco, December 1974, Abs.
- Scandone P., Giunta G., Liguori V., 1977. *The connection between the Apulia and Sahara continental margins in the Southern Apennines and in Sicily*. Mem. Soc. Geol. It., 13, p. 317-323.
- Schuster J.M., 1977. *Essai de reconstitution de l'histoire géologique et structurale de la Méditerranée Centrale*. Rev. I.F.P., v. 32, n. 4, p. 527-543.
- Semenza E., 1974. *La fase Giudicariense, nel quadro di una nuova ipotesi sull'orogenesi alpina nell'area italo-dinarica*. Mem. Soc. Geol. It., 13, p. 187-226.
- Selli R., ed., 1970. *Ricerche geologiche preliminari nel Mar Tirreno*. Giorn. Geol., v. 37, 249 pp.
- Selli R., Fabbri M., 1971. *Tyrrhenian: a Pliocene deep sea*. R.C. Accad. Lincei, v. 8, p. 104-116.
- Sikosek B., Maksimovic B., 1971. *Regional geotectonic pattern of the Adriatic sea belt*. Nafta, Zagreb, v. 4-5 (XXII), p. 298-304.
- Smith A.C., 1971. *Alpine deformation and the oceanic areas of the Tethys, Mediterranean and Atlantic*. Geol. Soc. Amer. Bull., v. 82, p. 2039-2070.
- Soffel H., 1972. *Anticlockwise rotation of Italy between Eocene and Miocene, paleomagnetic evidence*. Earth Planet. Sci. Lett., v. 17, p. 207-210.
- Stanley D.J., 1974a. *Basin plains in the eastern Mediterranean: significance in interpreting ancient marine deposits, II. Basin Distribution*. Bull. Cent. Rech. Pau-SNPA, v. 8, p. 373-388.
- Stride A.H., Belderson R.H., Kenyon N.H., 1977. *Evolving miogeanticlines of the Eastern Mediterranean (Hellenic, Calabrian and Cyprus Outer Ridges)*. Philosophical Transactions of the Royal Society of London, v. 284, p. 255-285.
- Subbotin S., Sollogub V., Prosen D., Dragesevic T., Mituch E., Posgay R., 1969. *Junction of deep structures of the Carpatho-Balkan region with those of the Black and Adriatic seas*. Can. J. Earth. Sci., v. 5, p. 1027-1035.
- Summerhayes C.P., Marks N., 1975. *Nile Delta: nature, evolution and collapse of continental shelf sediment system*. In: Proceedings UNESCO Seminar on Nile Delta Sedimentology, UNESCO Tech. Rept. 2, Alexandria, 29 pp.
- Tapponnier P., 1977. *Evolution tectonique du système alpin en Méditerranée: poinçonnement et écrasement rigide plastique*. Bull. Soc. Geol. Fr., (7), 19: 437-460.
- Vai G.B., 1979. *Tracing the Hercynian Structural Zones across "Neo-Europa": an Introduction*. Mem. Soc. Geol. It., v. 20, pp. 39-45.
- Van Bemmelen R.W., 1969. *Origin of the western Mediterranean Sea*. Verhandl. Kon. Ned. Geol. Mijnbouwk. Gen., v. 26, p. 13-52.
- Vandenberg, 1979. *The Tertiary movement of the Adriatic continental block*. Eos, 60, 604 (abs).
- Vezzani L., 1973. *L'Appennino siculo-calabro-lucano*. In: Moderne Vedute sulla Geologia dell'Appennino. Roma: Accad. naz. Lincei, p. 15-48.
- Villari L., 1969. *On particular ignimbrites of the island of Pantelleria (Channel of Sicily)*. Bull. Volcanol., v. 33, p. 1-12.
- Vogt P.R., Higgs R.H., 1969. *An aeromagnetic survey of the eastern Mediterranean Sea and its interpretation*. Earth Planet. Sci. Lett., v. 5, p. 439-448.
- Vogt P.R., Higgs R.H., Johnson G.L., 1971. *Hypotheses on the origin of the Mediterranean basin: magnetic data*. J. Geophys. Res., v. 76, p. 3207-3228.
- Wezel F.C., 1974. *Flysch successions and the tectonic evolution of Sicily during the Oligocene and Early*

- Miocene*. In: Geology of Italy, Squyres C., ed., Tripoli: Petrol., Expl. Soc., Libya, p. 1-23.
- Winnock E., 1979. *Les fosses du chenal de Sicilie*. In: The Pelagian Sea. Annales de l'Université de Provence, v. 6, p. 41-49.
- Winnock E., Bea F., 1979. *Structure de la mer Pelagienne*. In: Burolet et al., (eds.), 1979, p. 34-40.
- Wong H.K., Zarudzki E.F.K., Phillips J.D., Giermann G.K.F., 1971. *Some geophysical profiles in the eastern Mediterranean*. Bull. Geol. Soc. Amer., v. 82, p. 91-100.
- Woodside J.M., 1977. *Tectonic elements and crust in the eastern Mediterranean*. Mar. Geophysics Res., v. 3, p. 317-354.
- Yelnikov I.N., 1966. *Seismic reflection surveys in the Mediterranean between Crete and Africa*. Akad Nauk SSSR, Inst. Okeanol. (Transl.), v. 5.
- Yemel'yanov Ye M., Mikhaylov O.V., Shimkus K.M., Moskalenko V.N., 1968. *Some special features of the geomorphological structure and tectonic development of the Mediterranean Sea*. Acad. Sci. USSR, Oc. Comm. (Transl.), p. 34.
- Zammit Maempel G., 1977. *An Outline of Maltese Geology*. 1-44, Malta.
- Zarudzki E.F.K., 1972. *The Strait of Sicily - a geophysical study*. Rev. Géogr. Phys. Géol. Dyn., v. 14, p. 11-28.

BOOKS REVIEW

J.D. Byerlee and M. Wyss (Ed): *Rock friction and earthquake prediction*.
Contributions to Current Research in Geophysics (CCRG) 6. Birkhauses, Verlag, Basel und Stuttgart.

This is the special volume 116 n. 4-5 of Pure and Applied Geophysics and consists of 26 papers, most of which were presented at the Conference "Experimental studies of rock friction with application to earthquake prediction" held at Stanford University on April 28-30, 1977 and were integrated with suggestions arisen during the discussion at the congress.

The mechanics of the rocks under stress and the theory of dilatancy is of course the subject of the majority of papers and many experimental results obtained in laboratory are described.

Four articles at the beginning give a general description of the argument and they can be considered as reviews of selected topics.

A number of papers describes the relation between friction and rocks type, temperature, sliding rate, presence of water and the character of the sliding surface. Heat flow anomalies along active faults, such as the San Andreas fault, are compared in some articles with the laboratory measurements of friction and the formulated considerations are interesting. The temperature increase during frictional sliding is presented together with the results of some experiments.

The problem of the premonitory change in the seismic velocity through seismic regions is very important in earthquake prediction. The results obtained in California lead to some considerations about the precision required in the readings of the first arrivals. All the conditions of stress, pressure, temperature and pore pressure have to be investigated to study completely the problem of the velocity changes.

Other anomalous variations in physical properties reported sometimes before large earthquakes, such as in electrical resistivity, and in magnetic susceptibility as well as premonitory slip and gas emission have been studied in some laboratory experiments, and the results are shown.

The variations in dilatancy and permeability during changes in stress on rock were investigated in laboratory and some works are described.

Theoretical and experimental studies about particle and rupture velocity, modes of slip during fracture and frictional sliding and radiation of seismic waves are also presented in some papers.

The volume results in a rather complete presentation of the aspects connected with the subject of the rock friction.

Dario Slejko